

Jet-FTIR-spectroscopy of pyrrole clusters

Ingo Dauster, Corey A. Rice and Martin A. Suhm

Institute of Physical Chemistry, University Göttingen
Tammannstr. 6, 37077 Göttingen, Germany

61st Ohio State University International Symposium
on Molecular Spectroscopy
June 19–23, 2006

TB10 - Infrared / Raman

Outline

1 Introduction

- Motivation
- Experimental setup
- Theory
- Overview


2 Results

- Pyrrole as H-bond donor
- 2,5-Dimethylpyrrole as H-bond donor
- Conclusion

3 Pyrrole-2-carboxaldehyde

- Structure
- Filet-Jet spectra
- Structure of the dimer
- Frequency calculations

Motivation




Pyrrole



GEORG-AUGUST-UNIVERSITÄT
GÖTTINGEN

Motivation




Properties of pyrrole:

- **heterocyclic aromatic ring**
- electron rich aromatic (lone pair of N delocalized in aromatic ring)
- N–H functionality \Rightarrow H-bond donor
- π -electrons \Rightarrow H-bond acceptor (unconventional)
- \hookrightarrow self clustering (N–H \cdots π)



Motivation




Properties of pyrrole:

- heterocyclic aromatic ring
- **electron rich aromatic (lone pair of N delocalized in aromatic ring)**
- N–H functionality \Rightarrow H-bond donor
- π -electrons \Rightarrow H-bond acceptor (unconventional)
- \hookrightarrow self clustering (N–H \cdots π)



Motivation




Properties of pyrrole:

- heterocyclic aromatic ring
- electron rich aromatic (lone pair of N delocalized in aromatic ring)
- **N–H functionality \Rightarrow H-bond donor**
- π -electrons \Rightarrow H-bond acceptor (unconventional)
- \hookrightarrow self clustering (N–H \cdots π)



Motivation




Properties of pyrrole:

- heterocyclic aromatic ring
- electron rich aromatic (lone pair of N delocalized in aromatic ring)
- N–H functionality ⇒ H-bond donor
- π -electrons ⇒ H-bond acceptor (unconventional)
- \hookrightarrow self clustering (N–H \cdots π)



Motivation




Properties of pyrrole:

- heterocyclic aromatic ring
- electron rich aromatic (lone pair of N delocalized in aromatic ring)
- N–H functionality \Rightarrow H-bond donor
- π -electrons \Rightarrow H-bond acceptor (unconventional)
- ↗ self clustering (N–H \cdots π)




Motivation

Schematic structure of the pyrrole dimer:




Motivation



Porphyrin



Experimental setup



Filet-Jet

(fine but *lengthy*)

Slit nozzle


$600 \times 0.2 \text{ mm}^2$

C. A. Rice, N. Borho and M. A. Suhm, *Z. Phys. Chem.*, 219, 2005, 379–388.




GEORG-AUGUST-UNIVERSITÄT
GÖTTINGEN

Vibrational spectroscopy of clusters



Vibrational spectroscopy of clusters



X : π system,
O, etc.

red shift




Overview of the measured systems

H-bond donors



Pyrrole



2,5-Dimethylpyrrole


Pure H-bond acceptors



Benzene



N-Methylpyrrole



1,2,5-Trimethylpyrrole

Outline

1 Introduction

- Motivation
- Experimental setup
- Theory
- Overview


2 Results

- Pyrrole as H-bond donor
- 2,5-Dimethylpyrrole as H-bond donor
- Conclusion

3 Pyrrole-2-carboxaldehyde

- Structure
- Filet-Jet spectra
- Structure of the dimer
- Frequency calculations


Pyrrole



	$\tilde{\nu}_{\text{N-H}}/\text{cm}^{-1}$	$-\Delta\tilde{\nu}_{\text{N-H}}/\text{cm}^{-1}$
P	3533	—
P_2	3445	88
P_3	3393	140
P_4	3383	150




Pyrrole - structures of the clusters



P_2
Dimer



P_3
Trimer



P_4
Tetramer

B3LYP / 6-311++G**

A. Gómez-Zavaglia and R. Fausto, *J. Phys. Chem. A*, 108, **2004**, 6953–6967.



GEORG-AUGUST-UNIVERSITÄT
GÖTTINGEN

Comparison - experiment and DFT calculations


this work		Gómez-Zavaglia <i>et al.</i>	
in Helium		Calculation ^a	
	$\tilde{\nu}_{\text{N}-\text{H}}$ / cm ⁻¹	$-\Delta\tilde{\nu}_{\text{N}-\text{H}}$ / cm ⁻¹	$\tilde{\nu}_{\text{N}-\text{H}}^b$ / cm ⁻¹
P	3533	–	3593
P ₂	3445	88	3523
P ₃	3393	140	3495
P ₄	3383	150	3475

^a B3LYP / 6-311++G**


^b Scaling factor of 0.978

A. Gómez-Zavaglia and R. Fausto, *J. Phys. Chem. A*, 108, **2004**, 6953–6967.


Expansion in Helium vs. Argon



Expansion in Helium vs. Argon



Pyrrole in Helium and Argon




Comparision - Argon nanomatrix and Argon matrix

	this work		Gómez-Zavaglia <i>et al.</i>	
	in Argon		Argon matrix	
	$\tilde{\nu}_{\text{N-H}}$ / cm ⁻¹	$-\Delta\tilde{\nu}_{\text{N-H}}$ / cm ⁻¹	$\tilde{\nu}_{\text{N-H}}$ / cm ⁻¹	$-\Delta\tilde{\nu}_{\text{N-H}}$ / cm ⁻¹
P@Ar	3520	–	3523	–
P ₂ @Ar	3423	97	3418	105
P ₃ @Ar	3381	139	3396	127
P ₄ @Ar	–	–	3378	145

A. Gómez-Zavaglia and R. Fausto, *J. Phys. Chem. A*, 108, **2004**, 6953–6967.


Pyrrole / Benzene




	$\tilde{\nu}_{\text{N}-\text{H}} / \text{cm}^{-1}$	$-\Delta\tilde{\nu}_{\text{N}-\text{H}} / \text{cm}^{-1}$
P	3533	–
P+B	3485	48
P ₂	3445	88
P ₃	3393	140
P ₄	3383	150




Pyrrole as donor - overview




Pyrrole as donor - overview




Pyrrole as donor - overview




Pyrrole as donor - overview




Pyrrole as donor - overview



Pyrrole as donor - overview




2,5-Dimethylpyrrole




	$\tilde{\nu}_{\text{N}-\text{H}}/\text{cm}^{-1}$	$-\Delta\tilde{\nu}_{\text{N}-\text{H}}/\text{cm}^{-1}$
D	3506	—
—	3498	8
D_2	3420	86
D_3	3373	133




2,5-Dimethylpyrrole as donor - overview







2,5-Dimethylpyrrole as donor - overview






2,5-Dimethylpyrrole as donor - overview






Red-shift of the donor bands





Red-shift of the donor bands



Red-shift of the donor bands




Red-shift of the donor bands



⇒ universal order



Electron densities of the aromatic H-bond acceptors



Outline

1 Introduction

- Motivation
- Experimental setup
- Theory
- Overview

2 Results


- Pyrrole as H-bond donor
- 2,5-Dimethylpyrrole as H-bond donor
- Conclusion

3 Pyrrole-2-carboxaldehyde

- Structure
- Filet-Jet spectra
- Structure of the dimer
- Frequency calculations




Structure




GEORG-AUGUST-UNIVERSITÄT
GÖTTINGEN


Structure




Structure



Comparision with a peptide




Pyrrole-2-carboxaldehyde



Part of a peptide


Filet-Jet spectrum of the N–H stretching vibration




	$\tilde{\nu}_{\text{N-H}} / \text{cm}^{-1}$	$-\Delta\tilde{\nu}_{\text{N-H}} / \text{cm}^{-1}$
PCA	3493	–
(PCA) ₂	3273	220



Filet-Jet spectrum of the C=O stretching vibration




Structure of the dimer (top view)



B3LYP / 6-311++G**

GEORG-AUGUST-UNIVERSITÄT
GÖTTINGEN


Structure of the dimer (side view)



B3LYP / 6-311++G**


Frequency calculations

N-H stretching vibration




Frequency calculations

N–H stretching vibration




Frequency calculations

N–H stretching vibration




Frequency calculations

N–H stretching vibration




C=O stretching vibration




Frequency calculations

N–H stretching vibration




C=O stretching vibration




Frequency calculations

N–H stretching vibration




C=O stretching vibration




Frequency calculations

N–H stretching vibration



C=O stretching vibration



Acknowledgements



GEORG-AUGUST-UNIVERSITÄT
GÖTTINGEN

Acknowledgements



Acknowledgements



GEORG-AUGUST-UNIVERSITÄT
GÖTTINGEN

Acknowledgements



Financial support: GRK 782 of DFG
www.pcgg.de

GEORG-AUGUST-UNIVERSITÄT
GÖTTINGEN

Thank you
for your
attention!