

Manipulating the Motion of Large Molecules

Alternate Gradient Focusing and Deceleration of Benzonitrile



Jochen Küpper

*Fritz-Haber-Institut der MPG
Berlin*

21. June 2006



Cold Molecules



Applications of cold molecules

Experiments with molecules in high-field seeking states

- Getting complete control over both the internal and external degrees of freedom of molecules:
 - Beams of molecules in a single or small subset of quantum states
 - Spatially oriented molecules
 - Beams of molecules with computer-controlled velocities
 - Packets of trapped molecules
- Slow molecular beams for metrology, sensitive symmetry tests:
 - Weak interactions in chiral molecules
 - Time-reversal violating electric dipole moments of the electron (EDM)
 - Time-variation of fundamental constants (i.e. α/μ)
- Novel molecular beam collision, reaction, and interferometry experiments:
 - Conformational interchange and dynamics (*folding*)
 - Collisions at variable, well-defined energies
 - Scattering resonances
 - Quantum-controlled chemistry
- Ultra-low temperature phenomena:
 - Anisotropic dipole-dipole interaction:
 - repulsive: $\uparrow\uparrow$
 - attractive: $\rightarrow\rightarrow$
 - Molecular BEC


Applications of cold molecules

Experiments with molecules in high-field seeking states

- Getting complete control over both the internal and external degrees of freedom of molecules:
 - Beams of molecules in a single or small subset of quantum states
 - Spatially oriented molecules
 - Beams of molecules with computer-controlled velocities
 - Packets of trapped molecules
- Slow molecular beams for metrology; sensitive symmetry tests:
 - **Weak interactions in chiral molecules**
 - **Time-reversal violating electric dipole moment of the electron (EDM)**
 - Time-variation of fundamental constants (i. e. $d\alpha/dt$)
- Novel molecular beam collision, reaction, and interferometry experiments
 - **Conformational interchange and dynamics (*folding*)**
 - Collisions at variable, well-defined energies
 - Scattering resonances
 - **Quantum-controlled chemistry**
- Ultra-low temperature phenomena:
 - Anisotropic dipole-dipole interaction:
 - repulsive: $\uparrow\uparrow$
 - attractive: $\rightarrow\rightarrow$
 - Molecular BEC

Stark effect


Interaction of molecular charge distribution with external electric field:



Stark effect

Stark energy of $^{15}\text{ND}_3$


Interaction of molecular charge distribution with external electric field:



Stark effect of large molecules


Stark energy of benzonitrile (C_7H_5N)

Interaction of molecular charge distribution with external electric field:






Stark deceleration

Longitudinal motion








Deceleration of molecules in hfs states

Transverse motion (focusing)




Deceleration of molecules in hfs states

Transverse motion (focusing)




Longitudinal and transverse motion (deceleration and focusing)

independent control of transverse and longitudinal motion






Experimental setup

A modular Alternate Gradient decelerator




Experimental setup

A modular Alternate Gradient decelerator



Quantum state selective detection


Rotationally resolved laser induced fluorescence spectroscopy of benzonitrile ($S_1 \leftarrow S_0 0_0^0$)



inset: D.R. Borst, T.M. Korter, D.W. Pratt, *Chem. Phys. Lett.* **350**, 485 (2001)



Alternate Gradient focusing and guiding

Benzonitrile $S_0, v = 0, |J_{K_a K_c}\rangle = 0_{00}$





Alternate Gradient focusing and guiding

Benzonitrile $S_0, v = 0, |J_{K_a K_c}\rangle = 0_{00}$





Alternate Gradient focusing and guiding

Benzonitrile $S_0, v = 0, |J_{K_a K_c}\rangle = 0_{00}$




State selection

Guiding different eigenstates: $S_{0,v=0,|J_{KaKc}\rangle = 0_{00}$ vs 6_{06}



State selection


Guiding different eigenstates: Individual M -components of $S_0, v = 0, |J_{Ka}K_c\rangle = 6_{06}$



Decelerating ground-state benzonitrile

$$S_0, v = 0, |J_{K_a K_c}\rangle = 0_{00}$$



310 m/s \rightarrow 263 m/s: 28 % of kinetic energy is moved



Decelerating ground-state benzonitrile

$$S_0, v = 0, |J_{K_a K_c}\rangle = 0_{00}$$



310 m/s \rightarrow 263 m/s: 28 % of kinetic energy removed



Decelerating ground-state benzonitrile


$$S_0, v = 0, |J_{K_a K_c}\rangle = 0_{00}$$

310 m/s \rightarrow 263 m/s: 28 % of kinetic energy removed




Extended experimental setup – a modular decelerator

Decelerating large molecules to a *quasi*-standstill






Extended experimental setup – a modular decelerator




Decelerating large molecules to a *quasi*-standstill



AC traps for molecules in high-field seeking states








AC traps for molecules in high-field seeking states



van Veldhoven et al, *Phys. Rev. Lett.* **94**, 083001 (2005); Schnell et al, *in preparation*


AC traps for molecules in high-field seeking states



van Veldhoven et al, *Phys. Rev. Lett.* **94**, 083001 (2005); Schnell et al, *in preparation*

Decelerating larger molecules

Selecting tryptophan conformers



Conclusions

- Alternate Gradient focusing can be used to decelerate, bunch, focus, and trap polar neutral large and heavy molecules
 - Metastable CO in its hfs state has been decelerated from 250 m/s to 198 m/s
 - Benzonitrile (C_7H_5N) has been decelerated from 310 m/s to 263 m/s (removing $1/3$ of its kinetic energy)
 - ND_3 has been trapped in a hfs state
 - Alternate Gradient deceleration provides novel control of large and/or ground state molecules:
 - External motion can be precisely controlled
 - Internal state(s) can be selected from original sample
- Using the full experimental setup it will be possible to decelerate large molecules to a quasi-standstill in the laboratory frame and successively trap them
- Preparing intense beams of large, modular (bio-) molecules, i. e. using laser desorption, will allow novel (precision) studies of conformer-selected samples of such systems

Conclusions

- Alternate Gradient focusing can be used to decelerate, bunch, focus, and trap polar neutral large and heavy molecules
 - Metastable CO in its hfs state has been decelerated from 250 m/s to 198 m/s
 - Benzonitrile ($\text{C}_7\text{H}_5\text{N}$) has been decelerated from 310 m/s to 263 m/s (removing $\frac{1}{3}$ of its kinetic energy)
 - ND_3 has been trapped in a hfs state
 - Alternate Gradient deceleration provides novel control of large and/or ground state molecules:
 - External motion can be precisely controlled
 - Internal state(s) can be selected from original sample
- Using the full experimental setup it will be possible to decelerate large molecules to a quasi-standstill in the laboratory-frame and successively trap them
- Preparing intense beams of larger, modular (bio-) molecules, i.e. using laser desorption, will allow novel (precision) studies of conformer-selected samples of such systems

Conclusions

- Alternate Gradient focusing can be used to decelerate, bunch, focus, and trap polar neutral large and heavy molecules
 - Metastable CO in its hfs state has been decelerated from 250 m/s to 198 m/s
 - Benzonitrile ($\text{C}_7\text{H}_5\text{N}$) has been decelerated from 310 m/s to 263 m/s (removing $\frac{1}{3}$ of its kinetic energy)
 - ND_3 has been trapped in a hfs state
 - Alternate Gradient deceleration provides novel control of large and/or ground state molecules:
 - External motion can be precisely controlled
 - Internal state(s) can be selected from original sample
- Using the full experimental setup it will be possible to decelerate large molecules to a quasi-standstill in the laboratory-frame and successively trap them
- Preparing intense beams of larger, modular (bio-) molecules, i. e. using laser desorption, will allow novel (precision) studies of conformer-selected samples of such systems

Acknowledgment

Kirstin Wohlfart
Frank Filsinger

Hendrick L. Bethlem
Henrik Haak
Gerard Meijer

Melanie Schnell
Peter Lützow
Jacqueline van Veldhoven
Bretislav Friedrich



Max Planck Gesellschaft
Deutsche Forschungsgemeinschaft (SPP 1116)