

New Qualitative Features in Intramolecular Dynamics: Simple Examples

B. I. Zhilinskii^a, D. Sadovskii^a, K. Efstathiou^b

^a Université du Littoral, UMR 8101 du CNRS, 59 140

Dunkerque, France

^bRijksuniversiteit Groningen,


Groningen 9747 AC, The Netherlands

June 2007

62-th Symposium on Molecular Spectroscopy,

Columbus, Ohio

Monodromy of *swing-spring* with 1:1:2 resonance - model of Fermi resonance in CO₂



3D of freedom dynamical system - three resonant nonlinear oscillators - in the presence of axial symmetry.

CO₂ has four vibrational modes: symmetric and antisymmetric stretch and doubly degenerate bending. Antisymmetric vibration is out of resonance and can be “neglected” (averaged).

Integrable model

$$L = \frac{1}{2}(z_2\bar{z}_3 - \bar{z}_2z_3)i, \quad (1)$$

$$N = \bar{z}_1z_1 + \frac{1}{2}\bar{z}_2z_2 + \frac{1}{2}\bar{z}_3z_3, \quad (2)$$

$$H = aS + bR + cR^2 + E(N, L). \quad (3)$$

written in terms of invariant polynomials

$$R = \frac{1}{2}\bar{z}_2z_2 + \frac{1}{2}\bar{z}_3z_3 = (n_2 + n_3), \quad (4)$$

$$S = \frac{1}{4}(\bar{z}_1z_3^2 + z_1\bar{z}_3^2 + z_1\bar{z}_2^2 + \bar{z}_1z_2^2), \quad (5)$$

$$T = \frac{1}{4}(\bar{z}_1z_3^2 - z_1\bar{z}_3^2 - z_1\bar{z}_2^2 + \bar{z}_1z_2^2)i, \quad (6)$$

with $z = q - ip$, $\bar{z} = q + ip$, $\{z, \bar{z}\} = 2i$






Image of the energy momentum map for the $1 : 1 : 2$ resonant oscillator system with axial symmetry (*and without detuning*). Full 3D-image, typical constant- n section and fibers.







Matrix representation of monodromy for model with $1 : 1 : 2$ resonance

$$\begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (7)$$


The monodromy matrix is defined up to similarity transformation
 $M \sim AMA^{-1}$ with $A \in SL(3, R)$.




© Joyeux-Sadovskii 28-Jan-2004





Global view of the image of energy-momentum map for $L^2 = 0$ with small detuning.



Singular lines on the boundary of the image of the energy momentum map. The detailed representation of the boundary within the region near $E = x = \ell = 0$ indicated by a big circle is given in the next figure.









Hydrogen atom in external electric and magnetic fields


$$H = \frac{1}{2}\mathbf{P}^2 - \frac{1}{|\mathbf{Q}|} + F_e Q_2 + F_b Q_1 + \frac{G}{2}(Q_2 P_3 - Q_3 P_2) + \frac{G^2}{8}(Q_2^2 + Q_3^2) = E,$$






Structure of the 1:1 zone. Different dynamical strata of the zone (left) correspond to vertices of the genealogy graph (right).


path $A_2 A_{11} A''_1 A_0$ with fixed $a^2 = 0.45$




path $B_0''B_1''A_1''A_0$ for parallel fields near Z



Changes (of the constant n section) of the stratified images of the energy-momentum map for detuned 1:1 systems.



Contours in the stratified range of the energy-momentum map in the 1:1 zone
which encircle isolated critical values o and o' and o'' .



1:2 systems with fractional bidromy (left) and fractional monodromy (right).

References

- Phys. Rep. **341**, 11, 85, 173 (2001) [Reviews “for physicists”]
Proc. R. Soc. A **463** 1771-1790 (2007) *H atom in fields*
Phys. Rev. Lett. **93** 024302 (2004); J. Math. Phys. **45** 5076-5100 (2004);
Ann. Phys (N.Y.) **322** 164-200 (2007) *CO₂*
SIAM J. Appl. Dyn. Syst. **3** 261-351 (2004); Eur. Phys. J D **42** 61-72
(2007); *tetrahedral molecules A₄, AB₄*
Spectrochim. Acta A **61** 2867-2885 (2005) *O₃*
Chem. Phys. Lett. **89** 439 (2003); Phys.Rev. A **69** 032504 (2004)
LiCN, HCN