

Using Diffusion Monte Carlo to Probe Rotationally Excited States

Andrew S. Petit and Anne B. McCoy

The Ohio State University

Putting Diffusion Monte Carlo Onto a Carousel



Andrew S. Petit and Anne B. McCoy
The Ohio State University

Why Diffusion Monte Carlo (DMC)???

Highly fluxional molecules

Why Diffusion Monte Carlo (DMC)???

Highly fluxional molecules

Large amplitude nuclear motion

Why Diffusion Monte Carlo (DMC)???

Highly fluxional molecules

Large amplitude nuclear motion



Converged variational
calculations are very challenging

Why Diffusion Monte Carlo (DMC)???

Highly fluxional molecules

Large amplitude nuclear motion

Converged variational
calculations are very challenging






Why Diffusion Monte Carlo (DMC)???

Highly fluxional molecules

Large amplitude nuclear motion

Converged variational
calculations are very challenging



The General Diffusion Monte Carlo Methodology

$$i\hbar \frac{\partial \Psi}{\partial t} = \sum_{i=1}^N -\frac{\hbar^2}{2m_i} \nabla_i^2 \Psi + V\Psi$$

$$\Psi(t) = e^{\frac{itE_{ref}}{\hbar}} \sum_k c_k e^{\frac{-itE_k}{\hbar}} \phi_k$$

The General Diffusion Monte Carlo Methodology

$$i\hbar \frac{\partial \Psi}{\partial t} = \sum_{i=1}^N -\frac{\hbar^2}{2m_i} \nabla_i^2 \Psi + V\Psi$$

$$\Psi(t) = e^{\frac{itE_{ref}}{\hbar}} \sum_k c_k e^{\frac{-itE_k}{\hbar}} \phi_k$$

$$\tau = \frac{it}{\hbar}$$

The General Diffusion Monte Carlo Methodology

$$i\hbar \frac{\partial \Psi}{\partial t} = \sum_{i=1}^N -\frac{\hbar^2}{2m_i} \nabla_i^2 \Psi + V\Psi$$

$$\Psi(t) = e^{\frac{itE_{ref}}{\hbar}} \sum_k c_k e^{\frac{-itE_k}{\hbar}} \phi_k$$




$$\frac{\partial \Psi}{\partial \tau} = \sum_{i=1}^N \frac{\hbar^2}{2m_i} \nabla_i^2 \Psi - V\Psi$$

$$\boxed{\tau = \frac{it}{\hbar}}$$

The General Diffusion Monte Carlo Methodology

$$i\hbar \frac{\partial \Psi}{\partial t} = \sum_{i=1}^N -\frac{\hbar^2}{2m_i} \nabla_i^2 \Psi + V\Psi$$

$$\Psi(t) = e^{\frac{itE_{ref}}{\hbar}} \sum_k c_k e^{\frac{-itE_k}{\hbar}} \phi_k$$



$$\frac{\partial \Psi}{\partial \tau} = \sum_{i=1}^N \frac{\hbar^2}{2m_i} \nabla_i^2 \Psi - V\Psi$$

$$\frac{\partial \rho}{\partial t} = D \nabla^2 \rho - k\rho$$


Classical Diffusion Equation

$$\tau = \frac{it}{\hbar}$$

The General Diffusion Monte Carlo Methodology

$$i\hbar \frac{\partial \Psi}{\partial t} = \sum_{i=1}^N -\frac{\hbar^2}{2m_i} \nabla_i^2 \Psi + V\Psi$$

$$\Psi(t) = e^{\frac{itE_{ref}}{\hbar}} \sum_k c_k e^{\frac{-itE_k}{\hbar}} \phi_k$$



$$\frac{\partial \Psi}{\partial \tau} = \sum_{i=1}^N \frac{\hbar^2}{2m_i} \nabla_i^2 \Psi - V\Psi$$

$$\frac{\partial \rho}{\partial t} = D \nabla^2 \rho - k\rho$$

Classical Diffusion Equation


$$\tau = \frac{it}{\hbar}$$

Solution via statistical simulation!!!

The General Diffusion Monte Carlo Methodology

$$i\hbar \frac{\partial \Psi}{\partial t} = \sum_{i=1}^N -\frac{\hbar^2}{2m_i} \nabla_i^2 \Psi + V\Psi$$

$$\Psi(t) = e^{\frac{itE_{ref}}{\hbar}} \sum_k c_k e^{\frac{-itE_k}{\hbar}} \phi_k$$




$$\frac{\partial \Psi}{\partial \tau} = \sum_{i=1}^N \frac{\hbar^2}{2m_i} \nabla_i^2 \Psi - V\Psi$$

$$\boxed{\tau = \frac{it}{\hbar}}$$

The General Diffusion Monte Carlo Methodology


$$i\hbar \frac{\partial \Psi}{\partial t} = \sum_{i=1}^N -\frac{\hbar^2}{2m_i} \nabla_i^2 \Psi + V\Psi$$



$$\frac{\partial \Psi}{\partial \tau} = \sum_{i=1}^N \frac{\hbar^2}{2m_i} \nabla_i^2 \Psi - V\Psi$$

$$\boxed{\tau = \frac{it}{\hbar}}$$


$$\Psi(t) = e^{\frac{itE_{ref}}{\hbar}} \sum_k c_k e^{\frac{-itE_k}{\hbar}} \phi_k$$



$$\Psi(\tau) = \sum_k c_k e^{-(E_k - E_{ref})\tau} \phi_k$$

The General Diffusion Monte Carlo Methodology


$$i\hbar \frac{\partial \Psi}{\partial t} = \sum_{i=1}^N -\frac{\hbar^2}{2m_i} \nabla_i^2 \Psi + V\Psi$$



$$\frac{\partial \Psi}{\partial \tau} = \sum_{i=1}^N \frac{\hbar^2}{2m_i} \nabla_i^2 \Psi - V\Psi$$

$$\boxed{\tau = \frac{it}{\hbar}}$$

$$\Psi(t) = e^{\frac{itE_{ref}}{\hbar}} \sum_k c_k e^{\frac{-itE_k}{\hbar}} \phi_k$$




$$\Psi(\tau) = \sum_k c_k e^{-(E_k - E_{ref})\tau} \phi_k$$

If $E_{ref} = E_0$

The General Diffusion Monte Carlo Methodology


$$i\hbar \frac{\partial \Psi}{\partial t} = \sum_{i=1}^N -\frac{\hbar^2}{2m_i} \nabla_i^2 \Psi + V\Psi$$



$$\frac{\partial \Psi}{\partial \tau} = \sum_{i=1}^N \frac{\hbar^2}{2m_i} \nabla_i^2 \Psi - V\Psi$$

$$\boxed{\tau = \frac{it}{\hbar}}$$

$$\Psi(t) = e^{\frac{itE_{ref}}{\hbar}} \sum_k c_k e^{\frac{-itE_k}{\hbar}} \phi_k$$




$$\Psi(\tau) = \sum_k c_k e^{-(E_k - E_{ref})\tau} \phi_k$$

If $E_{ref} = E_0$

$$\lim_{\tau \rightarrow \infty} \Psi(\tau) = c_0 \phi_0$$

The General Diffusion Monte Carlo Methodology


$$i\hbar \frac{\partial \Psi}{\partial t} = \sum_{i=1}^N -\frac{\hbar^2}{2m_i} \nabla_i^2 \Psi + V\Psi$$



$$\frac{\partial \Psi}{\partial \tau} = \sum_{i=1}^N \frac{\hbar^2}{2m_i} \nabla_i^2 \Psi - V\Psi$$

$$\boxed{\tau = \frac{it}{\hbar}}$$

$$\Psi(t) = e^{\frac{itE_{ref}}{\hbar}} \sum_k c_k e^{\frac{-itE_k}{\hbar}} \phi_k$$



$$\Psi(\tau) = \sum_k c_k e^{-(E_k - E_{ref})\tau} \phi_k$$

If $E_{ref} = E_0$

Equilibrium



$$\lim_{\tau \rightarrow \infty} \Psi(\tau) = c_0 \phi_0$$



A Simple Game of Chance



Use an ensemble of M δ -functions (walkers)
to represent N particle wave function

Anderson, J.B. *J. Chem. Phys.* **1975**, 63, 1499-1503.

Anderson, J.B. *J. Chem. Phys.* **1976**, 65, 4121-4127.



A Simple Game of Chance



Use an ensemble of M δ -functions (walkers)
to represent N particle wave function

$$\Psi(\tau) \cong \sum_{i=1}^M \delta^{3N}(\vec{x} - \vec{x}_i(\tau))$$

Anderson, J.B. *J. Chem. Phys.* **1975**, 63, 1499-1503.

Anderson, J.B. *J. Chem. Phys.* **1976**, 65, 4121-4127.



A Simple Game of Chance



Use an ensemble of M δ -functions (walkers) to represent N particle wave function

$$\Psi(\tau) \cong \sum_{i=1}^M \delta^{3N}(\vec{x} - \vec{x}_i(\tau))$$

$$\Psi(\tau + \delta\tau) \cong e^{-(\hat{V} - E_{ref})\delta\tau} e^{-\hat{T}\delta\tau} \Psi(\tau)$$

Anderson, J.B. *J. Chem. Phys.* **1975**, 63, 1499-1503.

Anderson, J.B. *J. Chem. Phys.* **1976**, 65, 4121-4127.



A Simple Game of Chance



$$\Psi(\tau + \delta\tau) \cong e^{-(\hat{V} - E_{ref})\delta\tau} e^{-\hat{T}\delta\tau} \Psi(\tau)$$

Anderson, J.B. *J. Chem. Phys.* **1975**, 63, 1499-1503.

Anderson, J.B. *J. Chem. Phys.* **1976**, 65, 4121-4127.



A Simple Game of Chance



$$\Psi(\tau + \delta\tau) \cong e^{-(\hat{V} - E_{ref})\delta\tau} e^{-\hat{T}\delta\tau} \Psi(\tau)$$

Anderson, J.B. *J. Chem. Phys.* **1975**, 63, 1499-1503.

Anderson, J.B. *J. Chem. Phys.* **1976**, 65, 4121-4127.



A Simple Game of Chance



$$\Psi(\tau + \delta\tau) \cong e^{-(\hat{V} - E_{ref})\delta\tau} e^{-\hat{T}\delta\tau} \Psi(\tau)$$



Diffusion of walkers
in configuration space

Anderson, J.B. *J. Chem. Phys.* **1975**, 63, 1499-1503.
Anderson, J.B. *J. Chem. Phys.* **1976**, 65, 4121-4127.





A Simple Game of Chance



$$\Psi(\tau + \delta\tau) \cong e^{-(\hat{V} - E_{ref})\delta\tau} e^{-\hat{T}\delta\tau} \Psi(\tau)$$



Diffusion of walkers
in configuration space

Anderson, J.B. *J. Chem. Phys.* **1975**, 63, 1499-1503.
Anderson, J.B. *J. Chem. Phys.* **1976**, 65, 4121-4127.






A Simple Game of Chance



$$\Psi(\tau + \delta\tau) \cong e^{-(\hat{V} - E_{ref})\delta\tau} e^{-\hat{T}\delta\tau} \Psi(\tau)$$

Walker birth

Diffusion of walkers
in configuration space



Anderson, J.B. *J. Chem. Phys.* **1975**, *63*, 1499-1503.
Anderson, J.B. *J. Chem. Phys.* **1976**, *65*, 4121-4127.





A Simple Game of Chance




$$\Psi(\tau + \delta\tau) \cong e^{-(\hat{V} - E_{ref})\delta\tau} e^{-\hat{T}\delta\tau} \Psi(\tau)$$

Walker birth
and death



Diffusion of walkers
in configuration space



Anderson, J.B. *J. Chem. Phys.* **1975**, 63, 1499-1503.
Anderson, J.B. *J. Chem. Phys.* **1976**, 65, 4121-4127.






A Simple Game of Chance



$$\Psi(\tau + \delta\tau) \cong e^{-(\hat{V} - E_{ref})\delta\tau} e^{-\hat{T}\delta\tau} \Psi(\tau)$$

Walker birth
and death

Diffusion of walkers
in configuration space



Anderson, J.B. *J. Chem. Phys.* **1975**, *63*, 1499-1503.
Anderson, J.B. *J. Chem. Phys.* **1976**, *65*, 4121-4127.





A Simple Game of Chance




$$\Psi(\tau + \delta\tau) \cong e^{-(\hat{V} - E_{ref})\delta\tau} e^{-\hat{T}\delta\tau} \Psi(\tau)$$

Walker birth
and death

Diffusion of walkers
in configuration space

Reference energy
updated every time step




Anderson, J.B. *J. Chem. Phys.* **1975**, *63*, 1499-1503.

Anderson, J.B. *J. Chem. Phys.* **1976**, *65*, 4121-4127.




Evolution of E_{ref} with τ





Evolution of E_{ref} with τ



Where are we???

Where are we???



Where are we???



Calculate ground state energies

Where are we???



Calculate ground state energies

Obtain Monte Carlo sampling of
ground state wave function

Where are we???




Calculate ground state energies

Obtain Monte Carlo sampling of
ground state wave function

What About Excited States???


Fixed Node DMC

Assumes some knowledge
of nodal surfaces



Fixed Node DMC


Assumes some knowledge
of nodal surfaces



Fixed Node DMC

Assumes some knowledge
of nodal surfaces

Modify Hamiltonian with
infinite barrier placed
outside of nodal region




Fixed Node DMC

Assumes some knowledge
of nodal surfaces

Modify Hamiltonian with
infinite barrier placed
outside of nodal region

$$\hat{H} = \hat{H}_0 + \begin{cases} 0, & r > r_{node} \\ \infty, & r \leq r_{node} \end{cases}$$




Fixed Node DMC

Assumes some knowledge
of nodal surfaces

Modify Hamiltonian with
infinite potential placed
outside of nodal region

Lowest energy state
of new Hamiltonian:

$$\hat{H} = \hat{H}_0 + \begin{cases} 0, & r > r_{node} \\ \infty, & r \leq r_{node} \end{cases}$$




Fixed Node DMC

Assumes some knowledge
of nodal surfaces

Modify Hamiltonian with
infinite potential placed
outside of nodal region

DMC simulations
performed in all
nodal regions




Fixed Node DMC

Assumes some knowledge
of nodal surfaces

Modify Hamiltonian with
infinite potential placed
outside of nodal region

DMC simulations
performed in all
nodal regions

Average energies in
both nodal regions
must be equal



Nodal Surfaces of Rotationally Excited States

Two Assumptions:

Nodal Surfaces of Rotationally Excited States

Two Assumptions:

Rotational component of
wave function ONLY
depends on $\{\theta, \chi, \phi\}$

Nodal Surfaces of Rotationally Excited States

Two Assumptions:

Rotational component of
wave function ONLY
depends on $\{\theta, \chi, \phi\}$

$$\langle \theta, \chi, \phi | J, K, M = 0 \rangle_{\pm} \cong \frac{T_J^{K, \pm}(\theta, \chi)}{\sqrt{2\pi}} = \frac{(-1)^K Y_J^K(\theta, \chi) \pm Y_J^{-K}(\theta, \chi)}{\sqrt{4\pi(1 + \delta_{K0})}}$$

Nodal Surfaces of Rotationally Excited States

Two Assumptions:

Rotational component of
wave function ONLY
depends on $\{\theta, \chi, \phi\}$

$$\langle \theta, \chi, \phi | J, K, M = 0 \rangle_{\pm} \cong \frac{T_J^{K, \pm}(\theta, \chi)}{\sqrt{2\pi}} = \frac{(-1)^K Y_J^K(\theta, \chi) \pm Y_J^{-K}(\theta, \chi)}{\sqrt{4\pi(1 + \delta_{K0})}}$$

Nodal surfaces given by: $\theta = \theta_{node}$ OR $\chi = \chi_{node}$

Obtaining the Rotational Coordinates

Define body fixed frame

Obtaining the Rotational Coordinates

Define body fixed frame


Eckart frame to minimize
rotation-vibration coupling

Obtaining the Rotational Coordinates

Define body fixed frame

Eckart frame to minimize
rotation-vibration coupling

Rotational coordinates are
Euler angles connecting
space fixed and Eckart frames




Obtaining the Rotational Coordinates

Define body fixed frame

Eckart frame to minimize rotation-vibration coupling

Rotational coordinates are Euler angles connecting space fixed and Eckart frames



$$\begin{bmatrix} c\theta c\chi c\phi - s\chi s\phi & -c\theta s\chi c\phi - c\chi s\phi & s\theta c\phi \\ c\theta c\chi s\phi + s\chi c\phi & -c\theta s\chi s\phi + c\chi c\phi & s\theta s\phi \\ -s\theta c\chi & s\theta s\chi & c\theta \end{bmatrix}$$

Where are we now???

Where are we now???



Where are we now???



DMC method for calculating rotationally excited states


Where are we now???



DMC method for calculating rotationally excited states

How Well Does It Work???

Where are we now???



DMC method for calculating rotationally excited states

How Well Does It Work???

Converged variational calculations

Where are we now???




DMC method for calculating rotationally excited states

How Well Does It Work???

Converged variational calculations


Experiment

Comparing DMC and RVIB4 Energies




Huang, X.; Carter, S.; Bowman, J.M. *J. Chem. Phys.* **2003**, *118*, 5431-5441.

Comparing DMC to Experiment for H₃O⁺




Tang, J.; Oka, T. *J. Molec. Spect.* **1999**, 196, 120-130.

Comparing DMC to Experiment




Tang, J.; Oka, T. *J. Molec. Spect.* **1999**, *196*, 120-130.
Araki, M.; Ozeki, H.; Saito, S. *Mol. Phys.* **1999**, *97*, 177-183.

Comparing DMC to Experiment




Tang, J.; Oka, T. *J. Molec. Spect.* **1999**, *196*, 120-130.
Araki, M.; Ozeki, H.; Saito, S. *Mol. Phys.* **1999**, *97*, 177-183.


What Causes Deviations at Large J ???




What Causes Deviations at Large J ???




What Causes Deviations at Large J ???




What Causes Deviations at Large J ???




What Causes Deviations at Large J ???



What Causes Deviations at Large J ???



What Causes Deviations at Large J ???



Conclusions and Future Work

Conclusions and Future Work

Very successful treatment of states with $J \leq 2$

Conclusions and Future Work

Very successful treatment of states with $J \leq 2$

Study effects of rotational
excitation on geometric structure

Conclusions and Future Work

Very successful treatment of states with $J \leq 2$

Study effects of rotational
excitation on geometric structure


Some deviations at high J

Conclusions and Future Work

Very successful treatment of states with $J \leq 2$

Study effects of rotational
excitation on geometric structure

Some deviations at high J




Conclusions and Future Work

Very successful treatment of states with $J \leq 2$

Study effects of rotational
excitation on geometric structure

Some deviations at high J



Generalization of approach to asymmetric tops

Acknowledgements



Dr. Anne B. McCoy



Annie Lesiak






Charlotte E. Hinkle


Sara E. Ray

Samantha
Horvath


How Good Are Our Nodal Surfaces?



Probing Geometric Effects of Rotational Excitation with DMC




Probing Geometric Effects of Rotational Excitation with DMC



Probing Geometric Effects of Rotational Excitation with DMC

	H ₃ O ⁺	D ₃ O ⁺		
State	$\langle r_{OH} \rangle (\text{\AA})$	$\langle \theta_{HOH} \rangle (^{\circ})$	$\langle r_{OD} \rangle (\text{\AA})$	$\langle \theta_{DOD} \rangle (^{\circ})$
$ 0, 0, 0\rangle_+$	0.99564 ± 0.00026	113.062 ± 0.041	$0.99070 \pm .00020$	112.584 ± 0.031
$ 10, 10, 0\rangle_+$	0.00107 ± 0.00027	0.334 ± 0.043	0.00053 ± 0.00021	0.166 ± 0.032
$ 10, 10, 0\rangle_-$	0.00103 ± 0.00027	0.334 ± 0.043	0.00057 ± 0.00021	0.173 ± 0.032
$ 10, 0, 0\rangle$	$.00236 \pm 0.00029$	0.218 ± 0.045	0.00126 ± 0.00022	0.155 ± 0.034

Rotation about C axis:



Primarily affects molecular shape

Rotation about A OR B axes:

Larger impact on size



A Simple Game of Chance



$$|\Psi(\tau + \delta\tau)\rangle \cong e^{-(\hat{V} - E_{ref})\delta\tau} e^{-\hat{T}\delta\tau} |\Psi(\tau)\rangle$$

E_0 not known *a priori*

$$E_{ref}(\tau) = \boxed{\langle V(\tau) \rangle} - \alpha \left(\frac{\boxed{N(\tau)} - N(0)}{N(0)} \right)$$

Average potential
energy of walkers

Instantaneous
population
of walkers



A Simple Game of Chance



$$e^{-\hat{T}\delta\tau} \delta^{3N}(\vec{x} - \vec{x}_i) \propto e^{-\frac{m_i (\vec{x} - \vec{x}_i)^2}{2\delta\tau}}$$

Diffusion of Walkers

Random Displacements in *i*th
Coordinate Taken from Gaussian of Width

$$\sigma_i = \sqrt{\frac{\delta\tau}{m_i}}$$





A Simple Game of Chance



$$e^{-\hat{T}\delta\tau} \delta^{3N}(\vec{x} - \vec{x}_i) = K e^{-\frac{m_i (\vec{x} - \vec{x}_i)^2}{2\delta\tau}}$$

Diffusion of Walkers

Random Displacements in *i*th
Coordinate Taken from Gaussian of Width

$$\sigma_i = \sqrt{\frac{\delta\tau}{m_i}}$$





A Simple Game of Chance



We Already Have the Wave Function

$$\Psi(\vec{x}, \tau) \propto \sum_{i=1}^{N(\tau)} \delta^{3N}(\vec{x} - \vec{x}_i(\tau))$$

GOAL:

$$|\Psi(\vec{x}, \tau)|^2 \propto \sum_{i=1}^{N(\tau)} w_i(\tau) \delta^{3N}(\vec{x} - \vec{x}_i(\tau))$$



A Simple Game of Chance



IF


$$|\Psi(\vec{x}, \tau)|^2 \propto \sum_{i=1}^{N(\tau)} w_i(\tau) \delta^{3N}(\vec{x} - \vec{x}_i(\tau))$$

THEN

$$\langle A \rangle = \frac{\sum_{i=1}^{N(\tau)} w_i(\tau) A(\vec{x}_i)}{\sum_{i=1}^{N(\tau)} w_i(\tau)}$$

Descendent Weighting


Imaginary Time:
 τ



Descendent Weighting


Imaginary Time:

$$\tau + N_{step}$$



Descendent Weighting


Imaginary Time:
 τ



Descendent Weighting

Imaginary Time:


$$\tau + N_{step}$$




Descendent Weighting

Imaginary Time:


$$\tau + N_{step}$$




Projecting Wave Function onto Rotational Coordinates




Projecting Wave Function onto Rotational Coordinates




Monitoring Equilibration of Simulations



Monitoring Equilibration of Simulations



Monitoring Equilibration of Simulations



Only Observed for $J \leq 1$

Improvement to Projections Onto Rotational Coordinates
Yielded Better Agreement With RVIB4 Energies

Better Indication of Equilibration
Than Monitoring Energy

The Re-Crossing Correction



The Re-Crossing Correction



The Re-Crossing Correction

Probability of Walker Death Due to Node Re-Crossing:

$$P = \text{Exp} \left[\frac{-d_i(\tau) d_i(\tau + \delta\tau) m_i}{\delta\tau} \right]$$

Distance From *i*th
Nodal Surface

$$d_i(\tau) = \theta(\tau) - \theta_{node}$$

OR

$$d_i(\tau) = \chi(\tau) - \chi_{node}$$


Effective Mass Associated
With *i*th Nodal Surface



Effective Masses for Symmetric Tops

$$\theta = \theta_{node}$$

$$m_i = \sqrt{\left(\frac{I_A(\tau) + I_A(\tau + \delta\tau)}{2} \right) \left(\frac{I_B(\tau) + I_B(\tau + \delta\tau)}{2} \right)}$$



$$\chi = \chi_{node}$$

$$m_i = \frac{I_C(\tau) + I_C(\tau + \delta\tau)}{2}$$