

Measurement of rotational levels of the homonuclear helium dimer cation by extrapolation of Rydberg series

Daniel Sprecher, Jinjun Liu, Martin Schäfer,
Tobias Krähenmann, and Frédéric Merkt

Laboratory of Physical Chemistry, ETH Zurich, Switzerland

International Symposium on Molecular Spectroscopy
Ohio State University, June 21-25, 2010
RE03, June 24, 9:04am, 2015 McPherson Lab

Introduction: the He_2^+ molecule

- three-electron system, candidate for highly accurate *ab initio* calculations

Introduction: the He_2^+ molecule


- three-electron system, candidate for highly accurate *ab initio* calculations
- not much experimental data available:
 - 9 rovibrational $v=0 \rightarrow 1$ transitions in ${}^3\text{He} {}^4\text{He}^+$ (uncertainty 18 MHz) [1]
 - 7 rovibronic $X \rightarrow A$ ($v=22, 23 \rightarrow 0, 1$) transitions in ${}^4\text{He}_2^+$ (uncer. 0.2-2 MHz) [2]

[1] Yu and Wing PRL **59**, 2055 (1987)

[2] Carrington *et al.* JCP **102**, 5979 (1995)

Introduction: the He_2^+ molecule

- three-electron system, candidate for highly accurate *ab initio* calculations
- not much experimental data available:
 - 9 rovibrational $v=0 \rightarrow 1$ transitions in ${}^3\text{He} {}^4\text{He}^+$ (uncertainty 18 MHz) [1]
 - 7 rovibronic $X \rightarrow A$ ($v=22, 23 \rightarrow 0, 1$) transitions in ${}^4\text{He}_2^+$ (uncer. 0.2-2 MHz) [2]
- many studies on the Rydberg states of He_2 [M.L. Ginter *et al.* 1965-89]



Picture taken from:
[Buchenau *et al.* JCP **95**, 8134 (1991)]


[1] Yu and Wing PRL **59**, 2055 (1987)

[2] Carrington *et al.* JCP **102**, 5979 (1995)

Introduction: the He_2^+ molecule

- three-electron system, candidate for highly accurate *ab initio* calculations
- not much experimental data available:
 - 9 rovibrational $v=0 \rightarrow 1$ transitions in ${}^3\text{He} {}^4\text{He}^+$ (uncertainty 18 MHz) [1]
 - 7 rovibronic $X \rightarrow A$ ($v=22, 23 \rightarrow 0, 1$) transitions in ${}^4\text{He}_2^+$ (uncer. 0.2-2 MHz) [2]
- many studies on the Rydberg states of He_2 [M.L. Ginter *et al.* 1965-89]
- He_2 is a Rydberg molecule according to G. Herzberg [3]:

A molecule which has an essentially repulsive ground state and all excited states are Rydberg states




Picture taken from:
[Buchenau *et al.* JCP **95**, 8134 (1991)]

[1] Yu and Wing PRL **59**, 2055 (1987)


[2] Carrington *et al.* JCP **102**, 5979 (1995)

[3] Herzberg, Ann. Rev. Phys. Chem. **38**, 27 (1987)


Rydberg states of atoms and molecules



Rydberg states of atoms and molecules





Experimental




Experimental

Experimental
chamber





Experimental

Nozzle and discharge




Experimental chamber




Experimental



Nozzle and discharge



Experimental chamber




Solid-state laser system




Experimental



Nozzle and discharge



Experimental chamber



Solid-state laser system




- pulse length ~ 40 ns
- bandwidth ~ 20 MHz
- pulse energies


@875nm: ~ 15 mJ
 @438nm: ~ 2 mJ
 @292nm: ~ 150 μ J

Experimental


Nozzle and discharge




Experimental chamber



- field delay: 2 μ s
- field strength: 12.8 V/cm




Solid-state laser system




- pulse length \sim 40 ns
- bandwidth \sim 20 MHz
- pulse energies

@875nm: \sim 15 mJ
 @438nm: \sim 2 mJ
 @292nm: \sim 150 μ J


Overview Q-region




Resolved fine structure of the triplet a state




Resolved fine structure of the triplet a state




Resolved fine structure of the triplet a state




The $N^+=5$ Rydberg series



The $N^+=1,3$ Rydberg series



Overview over all observed states



Extrapolation of Rydberg series with multichannel quantum defect theory (MQDT)


$$E_{n,\ell} = E_i - \frac{hcR_M}{(n - \delta_\ell)^2}$$

quantum defects from MQDT
(adjusted to reproduce
positions of 600 $n < 25$ states)

[Ginter *et al.* JCP **81**, 6013 (1984)]

[Raunhardt *et al.* JCP **128**,
164310 (2008)]

Extrapolation of Rydberg series with multichannel quantum defect theory (MQDT)




$$E_{n,\ell} = E_i - \frac{hcR_M}{(n - \delta_\ell)^2}$$

quantum defects from MQDT
(adjusted to reproduce
positions of 600 $n < 25$ states)


[Ginter *et al.* JCP **81**, 6013 (1984)]

[Raunhardt *et al.* JCP **128**,
164310 (2008)]

Results of the extrapolation




Results of the extrapolation



all values in cm^{-1}			
N^+	Exp.	Calc. [1]	Δ
5	$198.3666(4)$	198.39	-0.02
3	$70.9395(4)$	70.95	-0.01

[1] Carrington *et al.* JCP **102**, 5979 (1995)]

Results of the extrapolation



all values in cm^{-1}			
N^+	Exp.	Calc. [1]	Δ
5	198.3666(4)	198.39	-0.02
3	70.9395(4)	70.95	-0.01

[1] Carrington *et al.* JCP **102**, 5979 (1995)]

Conclusions

- Transitions to triplet np Rydberg states (n up to 150) of He_2 have been resolved.
- The ionization energy of the $a\ ^3\Sigma_u^+$ state of He_2 was determined with an uncertainty of 0.006 cm^{-1} (180 MHz).
- The energy spacing between the first three rotational states of He_2^+ could be extracted with an accuracy of 0.0004 cm^{-1} (12 MHz).



Merkt group
ETH Zurich
Switzerland
(March 2009)






Merkt group
ETH Zurich
Switzerland
(March 2009)



Molecular Rydberg states and MQDT




Eigen quantum defect $\mu_{\ell,\ell'}^{S,\Lambda}$

Quantization condition:

$$\det \left| U_{i\alpha} \sin[\pi(\mu_\alpha + \nu_i)] \right| = 0$$

with $\nu_i = n - \delta_i$



Triplet pp eigen quantum defects have been adjusted to reproduce positions of almost 600 $n < 25$ states:

[D.S. Ginter *et al.* JCP **81**, 6013 (1984)]
[M. Raunhardt *et al.* JCP **128**, 164310 (2008)]

$$E_n = E_i - \frac{hcR_N}{(n - \delta)}$$

HeNe stabilized etalon

