

HIGH RESOLUTION FAR INFRARED FOURIER TRANSFORM SPECTROSCOPY OF THE NH₂ RADICAL

Marie-Aline MARTIN-DRUMEL,
Olivier PIRALI, Didier BALCON & Michel VERVLOET

Institut des Sciences Moléculaires d'Orsay (ISMO)
SOLEIL synchrotron - AILES beamline

66th International Symposium on Molecular Spectroscopy



ASTROPHYSICAL BACKGROUND



ESO/APEX & MSX/IPAC/NASA

- 1st detection towards SgrB2

VAN DISHOEK et al., *Astrophys. J.* **416** (1993)

ASTROPHYSICAL BACKGROUND



ESO/APEX & MSX/IPAC/NASA

- 1st detection towards SgrB2

VAN DISHOEK et al., *Astrophys. J.* **416** (1993)

- Recently detected by Herschel (HIFI) towards G10.6-0.4 (W31)

PERSSON et al., *A&A* **521** (2010)

STATE OF THE ART

NH₂: extensively studied from the sub-mm to the UV

STATE OF THE ART

NH₂: extensively studied from the sub-mm to the UV

- Sub-mm / THz / FIR

MÜLLER et al., *J. Mol. Spect.* **195** (1999)

STATE OF THE ART

NH₂: extensively studied from the sub-mm to the UV

- Sub-mm / THz / FIR

MÜLLER et al., *J. Mol. Spect.* **195** (1999)

- FIR Fourier transform spectroscopy (one study up to date):

MORINO & KAWAGUCHI et al., *J. Mol. Spect.* **182** (1997)

- R=0.007 cm⁻¹
- Absorption spectroscopy in the ground vibrational state
- 200 pure rotational transitions reported ($N_{max}=12$)

STATE OF THE ART

NH₂: extensively studied from the sub-mm to the UV

- Sub-mm / THz / FIR

MÜLLER et al., *J. Mol. Spect.* **195** (1999)

- FIR Fourier transform spectroscopy (one study up to date):

MORINO & KAWAGUCHI et al., *J. Mol. Spect.* **182** (1997)


- $R=0.007 \text{ cm}^{-1}$
- Absorption spectroscopy in the ground vibrational state
- 200 pure rotational transitions reported ($N_{\max}=12$)

OBJECTIVE

High resolution and high temperature

SPECTROSCOPY OF THE NH₂ RADICAL

Electronic: fundamental state $\tilde{X} \ ^2B_1$




SPECTROSCOPY OF THE NH₂ RADICAL


Electronic: fundamental state \tilde{X}^2B_1

Vibrational: $3N - 6 = 3$ vibration modes (ν_1, ν_2, ν_3)


SYMMETRIC STRETCHING



BENDING



ASYMMETRIC STRETCHING




SPECTROSCOPY OF THE NH₂ RADICAL


Electronic: fundamental state \tilde{X}^2B_1

Vibrational: $3N - 6 = 3$ vibration modes (ν_1, ν_2, ν_3)


SYMMETRIC STRETCHING



BENDING



ASYMMETRIC STRETCHING



Rotational: Asymmetric top rotor

- Hund's case (b)
- Rotational quantum numbers: N, K_a, K_c
- (000), (010): *b*-type transitions

ROTATIONAL STRUCTURE

$$\frac{N, K_a}{K_a=0 \dots N}$$

ROTATIONAL STRUCTURE

Asymmetric Splitting

$$\underline{K_c = N - K_a}$$


$$\frac{N, K_a}{K_a = 0 \dots N}$$

$$\underline{K_c = N + 1 - K_a}$$

$$\begin{aligned} K_a + K_c = \\ N \text{ or } N + 1 \end{aligned}$$


ROTATIONAL STRUCTURE

Asymmetric Splitting Fine Structure




ROTATIONAL STRUCTURE

Asymmetric Splitting Fine Structure Hyperfine Structure




ROTATIONAL STRUCTURE

Asymmetric Splitting Fine Structure Hyperfine Structure




EXPERIMENTAL SET-UPS

	Emission	Absorption
Interferometer	Bruker IFS 125	
Spectral range		15-700 cm ⁻¹
Precursor		NH ₃
Discharge	RF	DC
Power	1000 W	1-3 kV, 0.6-2 A
Pressure	~10 mbar	~0.1 mbar (+ 2 mbar He)
Light Source	plasma	SR
Resolution	4·10 ⁻³ cm ⁻¹	1·10 ⁻³ cm ⁻¹
Accuracy (l.p.)	< 5·10 ⁻⁴ cm ⁻¹ (12 MHz)	< 1·10 ⁻⁴ cm ⁻¹ (3 MHz)
Length	0.7 m	24 m
Rot. temp.	2500 K	300 K




EXPERIMENTAL SET-UPS

	Emission	Absorption
Interferometer		Bruker IFS 125
Spectral range		15-700 cm ⁻¹
Precursor		NH ₃
Discharge	RF	DC
Power	1000 W	1-3 kV, 0.6-2 A
Pressure	~10 mbar	~0.1 mbar (+ 2 mbar He)
Light Source	plasma	SR
Resolution	4·10 ⁻³ cm ⁻¹	1·10 ⁻³ cm ⁻¹
Accuracy (l.p.)	< 5·10 ⁻⁴ cm ⁻¹ (12 MHz)	< 1·10 ⁻⁴ cm ⁻¹ (3 MHz)
Length	0.7 m	24 m
Rot. temp.	2500 K	300 K




EXPERIMENTAL SET-UPS

	Emission	Absorption
Interferometer		Bruker IFS 125
Spectral range		15-700 cm ⁻¹
Precursor		NH ₃
Discharge	RF	DC
Power	1000 W	1-3 kV, 0.6-2 A
Pressure	~10 mbar	~0.1 mbar (+ 2 mbar He)
Light Source	plasma	SR
Resolution	4·10 ⁻³ cm ⁻¹	1·10 ⁻³ cm ⁻¹
Accuracy (l.p.)	< 5·10 ⁻⁴ cm ⁻¹ (12 MHz)	< 1·10 ⁻⁴ cm ⁻¹ (3 MHz)
Length	0.7 m	24 m
Rot. temp.	2500 K	300 K




EXPERIMENTAL SET-UPS

	Emission	Absorption
Interferometer	Bruker IFS 125	
Spectral range		15-700 cm ⁻¹
Precursor		NH ₃
Discharge	RF	DC
Power	1000 W	1-3 kV, 0.6-2 A
Pressure	~10 mbar	~0.1 mbar (+ 2 mbar He)
Light Source	plasma	SR
Resolution	4·10 ⁻³ cm ⁻¹	1·10 ⁻³ cm ⁻¹
Accuracy (l.p.)	< 5·10 ⁻⁴ cm ⁻¹ (12 MHz)	< 1·10 ⁻⁴ cm ⁻¹ (3 MHz)
Length	0.7 m	24 m
Rot. temp.	2500 K	300 K




EXPERIMENTAL SET-UPS

	Emission	Absorption
Interferometer		Bruker IFS 125
Spectral range		15-700 cm ⁻¹
Precursor		NH ₃
Discharge	RF	DC
Power	1000 W	1-3 kV, 0.6-2 A
Pressure	~10 mbar	~0.1 mbar (+ 2 mbar He)
Light Source	plasma	SR
Resolution	4·10 ⁻³ cm ⁻¹	1·10 ⁻³ cm ⁻¹
Accuracy (l.p.)	< 5·10 ⁻⁴ cm ⁻¹ (12 MHz)	< 1·10 ⁻⁴ cm ⁻¹ (3 MHz)
Length	0.7 m	24 m
Rot. temp.	2500 K	300 K




RESOLUTION OF THE ASYMMETRIC SPLITTING




★ Absorption: Resolution of the asymmetric splitting for several new lines

RESOLUTION OF THE FINE STRUCTURE




★ Absorption: Resolution of the fine structure for several new lines

RESOLUTION OF THE ¹⁴N HYPERFINE STRUCTURE



★ Absorption: Resolution of the hyperfine structure for several new lines

TRANSITION INVOLVING EXCITED ENERGY LEVELS



★ Emission: Pure rotational transitions within (000) and $\nu_2 = 1$

ANALYSIS (IN PROGRESS)

- Use of the SPFIT/SPCAT suite

PICKETT et al., *J. Mol. Spect.* (1991)

ANALYSIS (IN PROGRESS)

- Use of the SPFIT/SPCAT suite

PICKETT et al., *J. Mol. Spect.* (1991)

- Fit: pure rotational transitions within (000)

- 684 previously reported transitions (0-2 THz \sim 0-60 cm⁻¹)

MÜLLER et al., *J. Mol. Spect.* **195** (1999)

- \sim 700 transitions observed at Soleil (20-700 cm⁻¹)

- Watson's A-reduced rotational Hamiltonian (I' representation)

- IR RMS = $1.1 \cdot 10^{-3}$ cm⁻¹

- RMS ERROR = 1.58

FIT

FIT 1

1191 lines (507 SOLEIL)

$$N_{max} = 12 \quad (K_a = 3)$$

$$K_{amax} = 9 \quad (N = 9)$$

73 parameters

IR RMS=0.00029 cm⁻¹
RMS ERROR=1.66

FIT 2

1365 lines (681 SOLEIL)

$$N_{max} = 24 \quad (K_a = 2)$$

$$K_{amax} = 9 \quad (N = 13)$$

86 parameters

IR RMS=0.00075 cm⁻¹
RMS ERROR=1.49

FIT 3

1385 lines (701 SOLEIL)

$$N_{max} = 24 \quad (K_a = 2)$$

$$K_{amax} = 11 \quad (N = 13)$$

86 parameters

IR RMS=0.00112 cm⁻¹
RMS ERROR=1.58

FIT

FIT 1

1191 lines (507 SOLEIL)
 $N_{max} = 12$ ($K_a = 3$)
 $K_{amax} = 9$ ($N = 9$)
 73 parameters

IR RMS=0.00029 cm⁻¹
 RMS ERROR=1.66

FIT 2

1365 lines (681 SOLEIL)
 $N_{max} = 24$ ($K_a = 2$)
 $K_{amax} = 9$ ($N = 13$)
 86 parameters

IR RMS=0.00075 cm⁻¹
 RMS ERROR=1.49

FIT 3

1385 lines (701 SOLEIL)
 $N_{max} = 24$ ($K_a = 2$)
 $K_{amax} = 11$ ($N = 13$)
 86 parameters

IR RMS=0.00112 cm⁻¹
 RMS ERROR=1.58

- Light molecule: abnormal centrifugal distortion
- Renner-Teller effect

DRESSLER & RAMSAY, *Ph. Trans. Roy. Soc. London* **251** (1959)

- Unidentified transitions within (000) and (010)...

ROTATIONAL AND CENTRIFUGAL DISTORTION CONSTANTS (MHz)

Parameter	Present	Müller et al.
<i>A</i>	710 302.011 4(104)	710 302.131 9(250)
<i>B</i>	388 289.646 8(113)	388 288.871 8(1491)
<i>C</i>	245 013.373 9(70)	245 013.954 4(1316)
Δ_K	659.752 41(262)	659.765 23(1561)
Δ_{NK}	-125.142 86(169)	-125.045 19(1631)
Δ_N	31.686 89(50)	31.664 38(428)
δ_K	29.751 6(38)	29.375 5(761)
δ_N	12.709 480(182)	12.701 85(208)
$\Phi_K \cdot 10^3$	1 991.01(34)	1 991.49(350)
$\Phi_{KN} \cdot 10^3$	-283.00(46)	-267.88(472)
$\Phi_{NK} \cdot 10^3$	-49.337(121)	-48.98(139)
$\Phi_N \cdot 10^3$	13.069 4(128)	12.518 3(1195)
$\phi_K \cdot 10^3$	539.25(37)	490.64(778)
$\phi_{NK} \cdot 10^3$	-15.742(53)	-18.655(1194)
$\phi_N \cdot 10^3$	6.506 1(59)	6.294 3(432)
$L_K \cdot 10^6$	-10 626.3(162)	-10 156.(278)
$L_{NNK} \cdot 10^6$	63.68(139)	70.39(1281)
$L_N \cdot 10^6$	-9.846(135)	-4.786(792)
$L_{NK} \cdot 10^6$	-1 045.4(66)	-736.9(1070)
$L_{KKN} \cdot 10^6$	3 882.4(179)	2 472.(290)
$I_K \cdot 10^6$	-4 646.3(161)	-2 501.(208)
$I_{KN} \cdot 10^6$	72.0(32)	44.3(432)
$I_{NK} \cdot 10^6$	3.889(256)	45.87(818)
$I_N \cdot 10^6$	-4.891(67)	-3.043(295)
$P_K \cdot 10^6$	44.16(41)	49.01(426)
$S_K \cdot 10^6$	-0.119 4(38)	-0.184 4(299)

CONCLUSION

COMPLEMENT OF NH₂ DATABASE

~ 500 new transitions identified in (000)

CONCLUSION

COMPLEMENT OF NH₂ DATABASE

~ 500 new transitions identified in (000)

- **Emission:**

Observation of transitions involving excited energy levels

- (000) $N_{max} = 24$ ($K_a = 2$) et $K_{amax} = 12$ ($N = 24$)
- (010) (*in progress*)

CONCLUSION

COMPLEMENT OF NH₂ DATABASE

~ 500 new transitions identified in (000)

- **Emission:**

Observation of transitions involving excited energy levels

- (000) $N_{max} = 24$ ($K_a = 2$) et $K_{amax} = 12$ ($N = 24$)
- (010) (*in progress*)

- **Absorption:**

High resolution and accuracy on line position

- 1st resolution of the rotational splitting for ~ 40 transitions // (asymmetric splitting (10), fine (10) and hyperfine (20) structures)
- Improvement on line position accuracy for lines already reported ($< 3\text{MHz}$ / $\sim 1 \cdot 10^{-4}\text{cm}^{-1}$)

STUDY OF THE $^{15}\text{NH}_2$ ISOTOPE

Investigations on $^{14}\text{N}/^{15}\text{N}$ ratio

Objectives:

- Observation of pure rotational transitions of the $^{15}\text{NH}_2$ radical
- Identification on Herschel spectra ?

Collaboration:

- Astrophysicians: OBSPM, Evelyne ROUEFF; LRA, Maryvonne GERIN
- PhLAM: Laurent MARGULÈS, Stéphane BAILLEUX
- ISMO-SOLEIL: Olivier PIRALI, Marie-Aline MARTIN-DRUMEL

The THz Cosmos, June 24, Laurent MARGULÈS (FA06, 10h27)