

IMPROVED SPECTROSCOPIC ANALYSIS FOR THE $A\ ^3\Pi_{1u} - X\ ^1\Sigma_g^+$ SYSTEM OF Br₂

N. NISHIMIYA*, T. YUKIYA*,
and R. J. LE ROY**

*Department of Electronics and Information Technology, Tokyo Polytechnic University,
Iiyama 1583, Atsugi City, Kanagawa 243-0297, Japan

** Department of Chemistry, University of Waterloo, Waterloo, Ontario N2L 3G1,
Canada

Jun 20, 2011

1 Introduction

- Our research
- References

2 Energy Model of A – X System

3 Experimental Setup

4 Results

- Measuring Spectrum
- Assigned Band System
- Ro-vibrational Energy
- Spectroscopic Constants
- MLR Potential
- Direct Potential Fitting

5 Conclusion

Our Research

- $A - X$ system of $\text{ICl}^{(1)}$, $\text{IBr}^{(2),(4)}$, $\text{I}_2^{(3)}$ and $\text{Br}_2^{(5)}$ have been studied.
- In order to establish a frequency standard, the line positions of these spectra were determined.
- Spectroscopic constants of A and X -state were calculated.

(1) N. Nishimiya et al., "Nuclear Quadrupole Coupling Effect in the $A - X$ System and Spectroscopic Constants for $v = 2' - 4'$, $3' - 5'$ and $4' - 5'$ Progressions of ICl ", JMS, 163, 43(1994).

(2) Nishimiya et al., "Laser Spectroscopy of the $A^3\Pi_1 - X^1\Sigma^+$ System of IBr ", JMS, 173,8(1995).

(3) T. Yukiya et al., "High-Resolution Laser Spectroscopy of the $A^3\Pi_{1u} - X^1\Sigma_g^+$ System of I_2 with Ti: Sapphire Ring Laser", JMS, 182, 271(1997).

(4) T. Yukiya et al., "High-Resolution Laser Spectroscopy of the $A^3\Pi_1 - X^1\Sigma^+$ System of IBr with a Titanium:Sapphire Ring Laser", JMS, 214, 132(2002)

(5) Nishimiya et al., "High-Resolution Laser Spectroscopy of $A^3\Pi_{1u} - X^1\Sigma_g^+$ of Br_2 ", OHIO Columbus Meeting (2004).


References

- J.A.Horsley, "Rotational Analysis of the $A^3\Pi_{1u} - X^1\Sigma_g^+$ System of Bromine", J.Mol.Spectrosc., 22, 469 (1967)
- M.A.A.Clyne and J.A.Coxon, "The Emission Spectra of Br₂ and IBr Formed in Atomic Recombination Processes", J.Mol.Spectroscopy, 23, 58 (1967)
- J.A.Coxon "The $B3\Pi_{0u}^+ - X^1\Sigma_g^+$ system ⁷⁹Br⁷⁹Br., J.Mol.Spectrosc., 37, 39(1972)
- J.A.Coxon,et. al., "Absorption spectrum of the $A^3\Pi_{1u} - X^1\Sigma_g^+$ system of Br₂", J.Mol.Spectrosc., 41, 543 (1972)
- J.A.Coxon, "The Extreme Red Absorption Spectrum of Br₂, $A^3\Pi_{1u} - X^1\Sigma_g^+$, J.Mol.Spectrosc., 41, 548 (1972)
- J.A.Coxon, "Franck-Condon Factors and R-Centroids for Halogen Molecules-III.The $A^3\Pi_{1u} - X^1\Sigma_g^+$ System of ⁷⁹Br⁸¹Br", J.Mol.Spectroscopy, 41, 566 (1972).


References

- S. Gerstenkorn and P. Luc, "Analysis of the long range potential of ${}^{79}\text{Br}_2$ in the $B^3\Pi_{0u}^+$ state and molecular constants of the three isotopic bromine species ${}^{79}\text{Br}_2$, ${}^{79}\text{Br}{}^{81}\text{Br}_2$ and ${}^{81}\text{Br}_2$ ", J. Phys. France, 50, 1417 (1989)
- C. D. Boone, " Magnetic rotation study of the $A^3\Pi_{1u} - X^1\Sigma_g^+$ system of Br_2 ", Thesis (PhD), British Columbia (1999).
- C. Focsa, H. Li, and P. F. Bernath, "Characterization of the Ground State of Br_2 by Laser-Induced Fluorescence Fourier Transform Spectroscopy of the $B3\Pi_{0+u} - X^1\Sigma_g^+$ system", J.Mol.Spectroscopy, 200, 104 (2000).
- M. Saute, et. al., "Coefficients d'interaction à grande distance C_5 et C_6 pour les 23 états moléculaires de Cl_2 et de Br_2 ", Mol. Phys., 51, 1459 (1984)
- R. J. Le Roy, C. C. Haugen, J Tao, and H. Li "Long-range damping functions improve the short-range behaviour of 'MLR' potential energy functions", Mol. Phys., 109, 435 (2011)


Energy of A – X System



System Block Diagram



Assigned Line Quantum Number of A-state



Ro-vibrational Energy

The rotational and vibrational energies of lines are obtained by the equation:

$$\begin{aligned} \nu(v', J'; v'', J'') = & T_{v'} + B_{v'}^{e/f} \{J'(J' + 1) - \Lambda^2\} \\ & - D_{v'} \{J'(J' + 1) - \Lambda^2\}^2 \\ & + H_{v'} \{J'(J' + 1) - \Lambda^2\}^3 \\ & - \sum_{l=1} \sum_{m=0} Y''_{l,m} \left(v'' + \frac{1}{2}\right)^l \{J''(J'' + 1)\}^m \end{aligned} \quad (1)$$

Spectroscopic Constants

Spectroscopic Constants of $^{79,79}\text{Br}_2$ in the A-state

v'	T'_v	B'_v	$q'_v \times 10^5$	$D'_v \times 10^7$	$H'_v \times 10^{11}$
13	15453.760 (12)	0.045325 (42)		-1.33 (19)	-0.20 (23)
14	15521.1280(31)	0.043765 (12)		-1.560(67)	-0.190(95)
15	15581.5970(28)	0.0421330(86)		-1.731(45)	-0.240(60)
16	15635.6660(30)	0.0404762(55)		-1.837(24)	-0.340(28)
17	15684.0030(33)	0.0388558(59)		-1.966(26)	-0.370(31)
18	15727.3690(36)	0.0372876(63)		-2.063(27)	-0.410(31)
19	15766.4610(30)	0.0357611(55)	5.60(15)	-2.155(23)	-0.460(27)
20	15801.8810(25)	0.0343216(45)	5.60(27)	-2.235(20)	-0.550(24)
21	15834.0910(24)	0.0329272(46)	5.60(44)	-2.309(19)	-0.690(22)
22	15863.4350(25)	0.0315704(50)	5.90(38)	-2.432(22)	-0.830(27)
23	15890.1726(24)	0.0302285(48)	5.90(28)	-2.557(23)	-1.050(30)
24	15914.4858(25)	0.0288958(52)	5.80(23)	-2.734(28)	-1.310(40)

in cm^{-1} and σ in parentheses

Spectroscopic Constants

Spectroscopic Constants of $^{79,79}\text{Br}_2$ in the A-state - continued

v'	T'_v	B'_v	$q'_v \times 10^5$	$D'_v \times 10^7$	$H'_v \times 10^{11}$
25	15936.5210(48)	0.0275507(83)	5.60(34)	-2.938(43)	-1.660(65)
26	15956.3814(25)	0.0261880(63)	5.60(30)	-3.204(41)	-2.120(74)
27	15974.1603(20)	0.0247140(39)		-2.617(21)	-4.510(30)
28	15989.9010(35)	0.023364 (11)	5.50(100)	-3.930(81)	-3.60 (19)
29	16003.7040(37)	0.021876 (12)	5.70(54)	-4.36 (11)	-5.00 (27)
30	16015.6290(41)	0.020349 (16)		-4.78 (17)	-7.30 (51)
31	16025.7480(54)	0.018746 (31)		-5.51 (42)	-10.00 (160)
32	16034.1430(58)	0.017084 (35)		-6.34 (51)	-15.00 (210)
33	16040.9340(66)	0.015357 (56)		-7.50 (110)	-22.00 (630)
34	16046.2420(54)	0.013714 (34)		-13.30 (44)	
35	16050.236 (11)	0.011850 (80)		-16.00 (130)	
36	16053.0900(68)	0.009910 (85)		-19.00 (200)	
37	16054.987 (12)	0.00790 (31)		-20.0 (170)	

in cm^{-1} and σ in parentheses

Spectroscopic Constants

Spectroscopic Constants of $^{79,81}\text{Br}_2$ in the A-state

ν'	T'_{ν}	B'_{ν}	$q'_{\nu} \times 10^5$	$D'_{\nu} \times 10^7$	$H'_{\nu} \times 10^{11}$
2	14269.3310(38)	0.0560757(59)		-0.396 (24)	-0.010 (28)
3	14405.2524(23)	0.0553766(20)	1.270(41)	-0.4244(55)	-0.0120(41)
4	14536.1434(21)	0.0546292(15)	1.360(28)	-0.4546(42)	-0.0170(31)
5	14661.7889(21)	0.0538367(18)	1.500(30)	-0.4950(55)	-0.0210(47)
6	14781.9491(19)	0.0529907(13)	1.590(24)	-0.5430(37)	-0.0260(28)
7	14896.3779(19)	0.0520821(12)	1.710(21)	-0.5921(35)	-0.0420(27)
8	15004.8100(19)	0.0511055(13)	1.820(19)	-0.6645(38)	-0.0470(31)
9	15106.9630(19)	0.0500518(12)	2.010(18)	-0.7480(34)	-0.0650(25)
10	15202.5624(20)	0.0489104(15)	2.250(23)	-0.8518(46)	-0.0850(37)
11	15291.3646(21)	0.0476600(16)	2.560(22)	-0.9470(46)	-0.1350(36)
12	15373.1453(21)	0.0463185(18)	2.740(35)	-1.1010(56)	-0.1670(49)
13	15447.7965(24)	0.0448716(23)	3.220(34)	-1.2590(71)	-0.2200(62)
14	15515.3607(27)	0.0433337(29)	3.570(42)	-1.4290(96)	-0.2760(90)
15	15576.0740(40)	0.0417178(44)		-1.606 (14)	-0.320 (12)
16	15630.4050(39)	0.0400988(50)		-1.768 (16)	-0.350 (14)


in cm^{-1} and σ in parentheses

Dunham Coefficients for the X -state of ${}^{79,79}\text{Br}_2$



Y-parameters for the X -state	
$Y''_{1,0}$	$3.253167(11)\times 10^2$
$Y''_{2,0}$	-1.07937(34)
$Y''_{3,0}$	$-1.929(49)\times 10^{-3}$
$Y''_{4,0}$	$-1.83(36)\times 10^{-5}$
$Y''_{5,0}$	$-1.0(12)\times 10^{-7}$
$Y''_{6,0}$	$-3.0(16)\times 10^{-9}$
$Y''_{0,1}$	$8.21077(17)\times 10^{-2}$
$Y''_{1,1}$	$-3.198(10)\times 10^{-4}$
$Y''_{2,1}$	$-9.1(18)\times 10^{-7}$
$Y''_{3,1}$	$8.000(12)\times 10^{-10}$
$Y''_{4,1}$	$-4.0(26)\times 10^{-10}$
$Y''_{0,2}$	$-2.083(45)\times 10^{-8}$
$Y''_{1,2}$	$-1.7(28)\times 10^{-10}$
$Y''_{2,2}$	$1.3(50)\times 10^{-11}$
$Y''_{3,2}$	$-1.0(34)\times 10^{-12}$
$Y''_{4,2}$	$2.0(76)\times 10^{-14}$
$Y''_{0,3}$	$-4.0(320)\times 10^{-15}$

LeRoy-Bernstein Plot



LeRoy-Bernstein




Dependence of $B_{v'}$ and $\Delta B_{v'}$ on v' in $^{79,79}\text{Br}_2$ and $^{79,81}\text{Br}_2$



Dependence of $D_{v'}$ and $H_{v'}$ on v' in $^{79,79}\text{Br}_2$ and $^{79,81}\text{Br}_2$



Dependence of q_ν on ν in $^{79,79}\text{Br}_2$ and $^{79,81}\text{Br}_2$



Potential Equations

$$V_{MLR}(r) = \mathcal{D}_e \left\{ 1 - \frac{\mu_{LR}(r)}{\mu_{LR}(r_e)} e^{-\beta(r) \cdot y_p(r, r_e)} \right\}^2$$

$$\mu_{LR}(r) = C_{m1}/r^{m1} + C_{m2}/r^{m2} + \dots$$

$$\beta(r) = \beta_{MLR}(y_p(r, r_{ref})) = y_p(r, r_{ref})\beta_\infty + [1 - y_p(r, r_{ref})] \sum_{i=0}^{N_S, N_L} \beta_i y_p(r, r_{ref})^i$$

$$y_p = y_p(r, r_{ref}) = \frac{r^p - r_{ref}^p}{r^p + r_{ref}^p}$$

X-state

C5 cm ⁻¹ Å ⁵ :	-3.20	$\times 10^2$
C6 cm ⁻¹ Å ⁶ :	6.37	$\times 10^5$
C8 cm ⁻¹ Å ⁸ :	1.55	$\times 10^7$

A-state

C5 cm ⁻¹ Å ⁵ :	2.70	$\times 10^2$
C6 cm ⁻¹ Å ⁶ :	6.32	$\times 10^5$
C8 cm ⁻¹ Å ⁸ :	1.55	$\times 10^7$

All parameters are from M. Saute (Mol. Phys., 51, 1459(1984)).

Born–Oppenheimer Breakdown Correction Functions

$$\left\{ -\frac{\hbar^2}{2\mu_\alpha} \frac{d^2}{dr^2} + [V_{CN}(r) + \Delta V_{ad,W}^{(\alpha)}(r)] + \frac{\hbar^2[J(J+1) - \Lambda^2]}{2\mu_\alpha r^2} [1 + g_W^{(\alpha)}(r)] \right\} \psi_{v,J}(r) = E_{v,J} \psi_{v,J}(r)$$

$$\Delta V_{ad,W}^{(\alpha)}(r) = \frac{m_e}{M_A^\alpha} \tilde{S}_{ad,W}^A(r) + \frac{m_e}{M_B^\alpha} \tilde{S}_{ad,W}^B(r)$$

$$g_W^{(\alpha)}(r) = \frac{m_e}{M_A^\alpha} \tilde{R}_{na,W}^A(r) + \frac{m_e}{M_B^\alpha} \tilde{R}_{na,W}^B(r)$$

$$\tilde{S}_{ad,W}^A(r) = [1 - y_q(r, \textcolor{red}{r}_{ref})] \sum_{i=0} \mu_i^A y q_{ad}(r, r_{ref})^i + \mu_\infty^A y p_{ad}(r, r_{ref})$$

$$\tilde{R}_{na,W}^A(r) = [1 - y_p(r, \textcolor{red}{r}_{ref})] \sum_{i=0} \textcolor{red}{t}_i^A y q_{na}(r, \textcolor{red}{r}_e)^i + t_\infty^A y p_{na}(r, \textcolor{red}{r}_e)$$

Λ–Doubling for the A state


$$\left\{ -\frac{\hbar^2}{2\mu_\alpha} \frac{d^2}{dr^2} + [V_{ad}^1(r) + \Delta V_{ad}^{(\alpha)}(r)] + \frac{\hbar^2[J(J+1) - \Lambda^2]}{2\mu_\alpha r^2} [1 + g^{(\alpha)}(r)] \right.$$

$$\left. + s g_\Lambda(e/f) \Delta V_\Lambda^{(\alpha)}(r) [J(J+1)]^\Lambda \right\} \psi_{v,J}(r) = E_{v,J} \psi_{v,J}(r)$$


$$\Delta V_\Lambda^{(\alpha)}(r) = \left(\frac{\hbar^2}{2\mu_\alpha r^2} \right)^{2\Lambda} f_\Lambda(r)$$

$$f_\Lambda(r) = [1 - y_p(r, \mathbf{r}_e)] \sum_{i=0} \mathbf{w}_i^A y_p(r, \mathbf{r}_{r_e})^i + \mathbf{w}_\infty^A y_p(r, \mathbf{r}_e)$$


MLR Potentials for Br₂




Error Map of Calculated Line Positions without BOB



Error Map of Calculated Line Positions with BOB



DRMS Changing for Order of BOB



Parameters of MLR potential

	Without BOB	
	X-state	A-state
$D_e \text{ cm}^{-1}$	$1.6056930(17) \times 10^4$	$2.14814(23) \times 10^3$
$r_e \text{ \AA}$	2.2810513(20)	2.703341(46))
$C_5 \text{ cm}^{-1}\text{\AA}^5$	-3.30×10^2	3.20×10^4
$C_6 \text{ cm}^{-1}\text{\AA}^6$	6.274×10^5	6.370×10^5
$C_8 \text{ cm}^{-1}\text{\AA}^8$	1.550×10^7	1.550×10^7
β_0	0.791817	0.69237
β_1	0.8494	-0.6607
β_2	4.5933	0.1209
β_3	9.188	0.3429
β_4	10.84	-0.757
β_5	7.08	1.22
β_6	2.09	-5.20
β_7		2.85
β_9		-16.3
β_{10}		-42.2
β_{11}		22.0
β_{12}		44.6
β_{13}		-10.1
β_{14}		-20.2
β_∞	1.744098	0.3614602
$C_9(\text{eff})$	5.710675×10^5	5.710675×10^5
u_0	1.238	
	$N = 6, R_{ref} = 2.8$	$N = 14, p = q = 5, R_{ref} = 4.0$
	DMRS = 2.76032	

Parameters of MLR potential with BOB

		with BOB	
		X-state	A-state
T_e cm ⁻¹		1.39088290×10^4	
\mathcal{D}_e cm ⁻¹	$1.605683(39) \times 10^4$	2.14801×10^3	
r_e Å	$2.28102640(83)$	2.702	
C_5 cm ⁻¹ Å ⁵	-3.30×10^2	2.65×10^4	
C_6 cm ⁻¹ Å ⁶	6.274×10^5	6.370×10^5	
C_8 cm ⁻¹ Å ⁸	1.550×10^7	1.550×10^7	
β_0	0.791925	0.680218	t_0
β_1	0.85357	-0.68389	t_1
β_2	4.6088	0.0929	t_2
β_3	9.219	0.3091	t_3
β_4	10.96	-0.9675	t_4
β_5	7.25	1.299	t_5
β_6	2.000	-3.193	t_6
β_7		2.268	ω_0
β_8		12.127	ω_1
β_9		-16.27	ω_2
β_{10}		-18.456	ω_3
β_{11}		24.17	ω_∞
β_{12}		15.62	
β_{13}		-12.5	
β_{14}		-6.7	
β_∞	1.744052	0.36907150	
$C_9(eff)$	5.753939×10^5		
$C_{10}(eff)$		2.2449986×10^7	
$N = 6, R_{ref} = 2.8$		$N = 14, p = q = 5, R_{ref} = 4.0$	

Conclusion

- Spectra in the new bands were measured and assigned.
- The spectroscopic constants of each bands were determined with merged data.
- Dependence of B_v , q_v , D_v , H_v on v number was checked.
- BOB effect of line positions were verified.
- MLR potential was determined in the case with BOB and without BOB.