

# High Resolution Laser Spectroscopy of Rhodium Monobromide

A.G. Adam, T.F. Allen, L.E. Downie, and A.D.  
Granger

*Department of Chemistry and the Centre for Laser, Atomic, and Molecular Sciences, University of New Brunswick, Fredericton, NB, E3B 5A3*


C. Linton and D.W. Tokaryk

*Department of Physics and the Centre for Laser, Atomic, and Molecular Sciences, University of New Brunswick, Fredericton, NB, E3B 5A3*

## Previous Work on Pertinent Rhodium Diatomics


- **RhH**: Balfour et al., J. Mol. Spectroc. 201, 244(2000).
  - ${}^3\Delta_3$  from  $\delta^3\Pi^4\sigma^1$  configuration
- **RhF**: Li et al., J. Chem. Phys. 121, 2591(2004).
  - ${}^3\Pi_2$  from  $\delta^4\Pi^3\sigma^1$  configuration
- **RhCl**: Shepard et al., J. Mol. Spectrosc. 234, 99(2005).
  - ${}^3\Pi_2$  from  $\delta^4\Pi^3\sigma^1$  configuration
- **RhBr Theory**: Cheng et al., J. Comp. Chem. 28, 2190(2007).
  - ${}^3\Delta_3$  from  $\delta^3\Pi^4\sigma^1$  configuration for RhF, RhCl, RhI
  - ${}^3\Pi_2$  from  $\delta^4\Pi^3\sigma^1$  configuration for RhBr

# Apparatus





---

## Low Resolution Survey



# Dispersed Fluorescence



# Equilibrium Molecular Constants and Bond Strengths

	<i>Ground State X</i>	<i>Lower State X1</i>
$\omega_e$ (cm <sup>-1</sup> )	260(8)	268(8)
$\omega_{eX_e}$ (cm <sup>-1</sup> )	2.0(1)	6.0(2)
k (N/m)	180	190

RhF: 575 cm<sup>-1</sup>, 312 N/m

RhCl: 348 cm<sup>-1</sup>, 186 N/m


HF: 914 N/m

HCl: 516 N/m

HBr: 411 N/m


High Resolution

Strong Q-branch




High Resolution

Weak Q-branch




High Resolution

P-head




High Resolution

R-head




## Two Isotopologues



## $\Lambda$ -Doubling

Rh<sup>79</sup>Br



# Preliminary Polynomial Fits


Rh<sup>81</sup>Br

$\Omega' - \Omega''$	$T'_0$ (cm <sup>-1</sup> )	$B'$ (cm <sup>-1</sup> )	$D'$ (cm <sup>-1</sup> )	$B''$ (cm <sup>-1</sup> )	$D''$ (cm <sup>-1</sup> )	$q''$ (cm <sup>-1</sup> )	$q''_d$ (cm <sup>-1</sup> )
2 - 2	23612.37(1)				Perturbed		
2 - 2	23610.4944(8)	0.063003(5)	-1.09(2)x10 <sup>-6</sup>	0.064849(5)	3.79(11)x10 <sup>-9</sup>		
2 - 2	23369.0473(5)	0.064610(12)	4(5)x10 <sup>-9</sup>	0.064926(13)	2.13(56)x10 <sup>-8</sup>	4.54(47)x10 <sup>-9</sup>	
2 - 2	21934.5344(5)	0.065964(15)	5.77(82)x10 <sup>-8</sup>	0.064833(15)	6.89(85)x10 <sup>-8</sup>	4.72(81)x10 <sup>-9</sup>	
2 - 2	20641.0323(5)	0.064568(11)	-1.64(54)x10 <sup>-8</sup>	0.064953(11)	1(5)x10 <sup>-9</sup>		7.17(49)x10 <sup>-9</sup>


## Future Work

- Assign the remaining bands
- Perform a global fit
- Try to resolve the apparent hyperfine structure

R(2)



R(19)



## Acknowledgements

- Joyce MacGregor, Michelle Jacobs, Jessica Prescott, Rhys Ems, and Dr. Tim Steimle

## Funding



**NSERC  
CRSNG**



Questions?