

The visible spectrum of iridium monohydride and monodeuteride

A. G. ADAM AND A. D. GRANGER, Chemistry Department, and Centre for Lasers, and Atomic, and Molecular Sciences, University of New Brunswick, Fredericton, NB, E3B 5A3

C. LINTON AND D. W. TOKARYK, Physics Department, and Centre for Lasers, and Atomic, and Molecular Sciences, University of New Brunswick, Fredericton, NB, E3B 5A3


AND TIMOTHY C. STEIMLE, Department of Chemistry and Biochemistry, Arizona State University, Tempe, AZ 85287



June 22, 2011



Why should we study diatomic species?



Change in electronic character as a function of changing ligand or transition metal

Comparisons between theory and experiment

Transition metal complexes important catalytically

Iridium monohydride and monodeuteride - IrH/IrD


Motivations

- Prior work – no experimental studies
- Only five iridium-containing diatomics published to date
- Continues work on group 9-monohydrides (CoH, RhH, and IrH)


Synthesis of Diatomic Species

- Transition-metal source (metal rod)
- Reactant Source (reactant gas)
- Heat (UV laser)





Low resolution (IrD)




Determination of the ground state of IrH/IrD


- First lines R(4), Q(4), P(5)
- CoH (${}^3\Phi_i$) [Bernath (1996)], CoF (${}^3\Phi_i$) [Adam (1994)],, IrF (${}^3\Phi_i$) [Adam (2009)],
- Theory predicts either ${}^3\Delta$ [Cassarubios (1999), Wittborn (1995)] or ${}^3\Sigma^-$ [Balasubramanian (1991)]
- All studies predict a low-lying ${}^3\Phi$ state

$${}^3\Phi_4$$

Low resolution (IrH) with 'extra' lines



Dispersed fluorescence of 553 nm band of IrH



Vibrational Analysis

	Vibrational frequency (cm ⁻¹)	Anharmonicity term (cm ⁻¹)
IrH G.S.	2140(11)	--
IrH low-lying	~ 2100	--
IrD G.S.	1609(2)	19.3(5)
IrD low-lying	~ 2100	--

Rotational fit results

Molecule	Assignment	T (cm ⁻¹)	B (cm ⁻¹)	D (cm ⁻¹) x 10 ⁻³
IrH	$X^3\Phi_4$ v=0	0 fixed	6.540(4)	0.25 fixed
		18060.3(3)	6.67(3)	6.1(7)
		18459.2(3)	6.31(3)	4.4(7)
		19720.7(3)	6.89(3)	12.7(6)
		20488.3(2)	6.24(6)	--
		21189.9(3)	6.33(3)	11.1(7)
		21873.5(2)	5.88(1)	--
		22555.4(3)	6.17(3)	8.4(7)
		23466.4(2)	6.57(1)	--


Comparison with RhH and CoH

Parameter	B (cm ⁻¹)	r (Å)	ω_e (cm ⁻¹)	k (N/m)
CoH ¹	7.14	1.53	1859	202
RhH ²	6.55	1.59	2040	245
IrH	6.54	1.60	2140	271


¹Bernath 1996

²Balfour 2000

High resolution at UNB, Q(4) line



High resolution at Arizona State University (Steimle Lab), Q(4) line



Acknowledgements

UNB Spectroscopy Group

Allan Adam
Dennis Tokaryk
Colan Linton
Joyce MacGregor

Theoretical Work

Fritz Grein

Hyperfine Work (ASU)

Tim Steimle
Sarah Frey
Fang Wang
Anh Le

Funding

UNB
NSERC
ASU

Questions?