



UNB

Far-infrared synchrotron-based spectroscopy of furan: analysis of the ν_{14} - ν_{11} perturbation and the ν_{18} and ν_{19} levels




Canadian Centre canadien
Light de rayonnement
Source synchrotron

Dennis Tokaryk
**Dept. of Physics and Centre for Laser,
Atomic, and Molecular Sciences, UNB**



Canadian Centre canadien
Light de rayonnement
Source synchrotron




Scott Culligan, Dept. of Physics, UNB (now Oxford U.)
Dr. Brant Billinghamurst, Canadian Light Source
Dr. J. A. van Wijngaarden, Dept. of Chemistry, U. Manitoba



Bruker IFS 125HR high-resolution Fourier transform spectrometer

- 10 m path difference
- resolution of 0.001 cm^{-1}





With successive improvements, the synchrotron now has a significant advantage over the globar from $350 \sim 800 \text{ cm}^{-1}$, and reasonable advantage over the Hg lamp from $150\text{-}350 \text{ cm}^{-1}$.



UNB

Bruker IFS 125HR high-resolution Fourier transform spectrometer

- 10 m path difference
- resolution of 0.001 cm^{-1}



66th OSU International Symposium on Molecular Spectroscopy 2011




UNB


Simple 5-membered rings are large enough to have dense, complex spectra, but are easy to handle.

The molecules often absorb strongly just where we are seeing the greatest synchrotron advantage.


We have worked on pyrrole, furan, thiophene, pyrazole, and imidazole. They have been a great introduction to asymmetric rotor physics!




Furan




Pyrrole



Thiophene



Pyrazole





Imidazole



UNB

Furan: overview of our spectrum





Furan



UNB

Furan: overview of our spectrum





Furan



UNB




Furan: overview of our spectrum



Furan




Furan: overview of our spectrum







Furan





UNB














UNB


Furan: vibrational energy level structure






UNB

Furan: vibrational energy level structure






Furan: vibrational energy level structure






Furan: vibrational energy level structure






Furan: vibrational energy level structure





UNB


Furan: the $\nu_{18} - \nu_{11}$ transition





P(16)


R(16)





UNB

Furan: Reduced energies of the ν_{11} and ν_{14} levels





UNB


Fit to a Watsonian, A -reduction, III^r representation. All values in cm^{-1} .

| | Ground state | ν_{14} | ν_{11} | ν_{18} | ν_{19} |
|-------------------------|------------------|-----------------|-----------------|-----------------|-----------------|
| T_0 | | 602.8445069(94) | 599.693665(26) | 1266.750171(15) | 1180.836732(13) |
| A | 0.3151220975(74) | 0.314310966(43) | 0.31432994(30) | 0.31529597(41) | 0.31464163(24) |
| B | 0.3084381338(75) | 0.307602792(42) | 0.30779360(29) | 0.30884839(38) | 0.30803831(19) |
| C | 0.1558019445(69) | 0.155846701(23) | 0.155870786(60) | 0.155686313(35) | 0.155512886(30) |
| $D_J \times 10^8$ | 11.11682(66) | 10.99943(92) | 11.1629(40) | 11.2596(44) | 11.1412(31) |
| $D_{JK} \times 10^8$ | -17.5584(11) | -17.0619(16) | -17.454(11) | -17.8158(84) | -17.6731(60) |
| $D_K \times 10^8$ | 7.6175(12) | 7.2489(18) | 7.4746(96) | 7.7242(47) | 7.6898(37) |
| $d_1 \times 10^8$ | -0.17487(57) | -0.2147(11) | -0.5716(60) | -0.797(21) | -0.8608(76) |
| $d_2 \times 10^8$ | 0.1440(11) | 0.1582(12) | 0.1429(31) | 0.1711(89) | 0.1737(39) |
| $H_{JK} \times 10^{13}$ | -1.738(46) | | | | |
| $H_{KJ} \times 10^{13}$ | 1.746(60) | | | | |
| $C_{Cor} \times 10^4$ | | 1.15332(44) | | | |

RMSE: 0.0001 cm^{-1}




UNB





UNB

Possible cause of the second-order Coriolis resonance?





What we've learned so far...

- Perturbations in the ν_{11} and ν_{14} levels of furan are the result of a type-c Coriolis interaction.
- The second-order interaction with $\Delta K_c = \pm 2$ is the most obvious due to an accidental near-degeneracy of $K_c = 6$ lines in the ν_{14} state with $K_c = 4$ levels in the ν_{11} state.
- This analysis would have been extremely difficult without the advantage provided by far-IR synchrotron radiation coupled to a FT spectrometer run at very high resolution.




UNB

Bruker IFS 125HR high-resolution Fourier transform spectrometer

- 10 m path difference
- resolution of 0.001 cm^{-1}

Synchrotron details:

- injections every 8 hours
- max. current of 250 mA, drops to $\sim 150 \text{ mA}$ before the next injection.



We run helium-cooled Cu:Ge and Si:bolometer detectors, as well as nitrogen-cooled MCTs

With successive improvements, the synchrotron now has a significant advantage over the globar from $350 \sim 800 \text{ cm}^{-1}$, and reasonable advantage over the Hg lamp from $150\text{-}350 \text{ cm}^{-1}$.