


Quantum state-resolved reactive and inelastic scattering at gas-liquid and gas-solid interfaces

Monika Grütter, Daniel J. Nelson, and David J. Nesbitt

JILA, University of Colorado and NIST, and Department of Chemistry
and Biochemistry, University of Colorado, Boulder, CO 80309




FONDS NATIONAL SUISSE
SCHWEIZERISCHER NATIONALFONDS
FONDO NAZIONALE SVIZZERO
SWISS NATIONAL SCIENCE FOUNDATION

Motivation

Scattering of gas beams from liquid and solid surfaces:


inelastic scattering



→ heterogeneous chemistry
aerosols
heterogeneous catalysis

→ dynamics of energy transfer at interfaces
→ quantum-state resolution: observe all internal degrees of freedom
(translational, rotational, vibrational)

reactive scattering



A. M. Zolot et al., JCP, 129, 194705, 2008


Experimental overview

How to obtain quantum-state resolution in a scattering process:

- gas: supersonic expansion, skimmed
 - cooling of internal degrees of freedom
 - control over collision energy
- heated gold surface (500 K): locally atomically flat (111) surface
 - no reaction dynamics
- Resonance-enhance multi-photon ionization (REMPI)
 - vibrational and rotational resolution
- Velocity-map imaging (VMI)
 - translational degree of freedom

Two main channels in a scattering process

Former work: HCl inelastically scattered from a heated gold surface [1]:



Velocity map image integrated over all rotational states of HCl:

TD: Trapping-desorption

- equilibrium with surface
- isotropic velocity distribution


IS: Impulsive scattering

- hyperthermal
- forward-scattered

[1] J.R.Roscioli, D.J.Bell, D.J.Nelson and D.J.Nesbitt,
PhysChemChemPhys, **14**, 4070, 2012

Internal degrees of freedom


Information on rotational and translational energy distributions [1]:



[1] J.R.Roscioli, D.J.Bell, D.J.Nelson and D.J.Nesbitt,
PhysChemChemPhys, **14**, 4070, 2012

Internal degrees of freedom

Information on rotational and translational energy distributions [1]:




[1] J.R.Roscioli, D.J.Bell, D.J.Nelson and D.J.Nesbitt,
PhysChemChemPhys, **14**, 4070, 2012

Vibrational excitation in inelastic scattering

HCl:
diatomic molecule

$$\omega_e \sim 2890 \text{ cm}^{-1}$$

No observable collision
energy transfer into vibration



Benzene:
symmetric top of D_{6h} symmetry
20 normal modes
prototype hydrocarbons

- $\omega_{20} \sim 405 \text{ cm}^{-1}$ C-C-C bend
- $\omega_{19} \sim 970 \text{ cm}^{-1}$ C-H bend
- $\omega_{11} \sim 845 \text{ cm}^{-1}$ C-H bend
- $\omega_8 \sim 703 \text{ cm}^{-1}$ C-C-C bend
- $\omega_7 \sim 992 \text{ cm}^{-1}$ C-H bend
- $\omega_6 \sim 608 \text{ cm}^{-1}$ C-C-C bend
- $\omega_4 \sim 673 \text{ cm}^{-1}$ C-H bend
- $\omega_2 \sim 993 \text{ cm}^{-1}$ C-C stretch


- [2] G. Herzberg, *Mol. Spectra & Mol. Structure, Volume III*, Krieger, 1991
- [3] E. Riedle et al., *JCP*, **91**, 4555, 1989
- [4] M. Okruss et al., *JMS*, **193**, 293, 1999

Vibrational excitation in inelastic scattering

HCl:
diatomic molecule

$$\omega_e \sim 2890 \text{ cm}^{-1}$$

No observable collision
energy transfer into vibration



Benzene:
symmetric top of D_{6h} symmetry
20 normal modes
prototype hydrocarbon

- $\omega_{20} \sim 405 \text{ cm}^{-1}$ C-C-C bend
- $\omega_{19} \sim 970 \text{ cm}^{-1}$ C-H bend
- $\omega_{11} \sim 845 \text{ cm}^{-1}$ C-H bend
- $\omega_8 \sim 703 \text{ cm}^{-1}$ C-C-C bend
- $\omega_7 \sim 992 \text{ cm}^{-1}$ C-H bend
- $\omega_6 \sim 608 \text{ cm}^{-1}$ C-C-C bend
- $\omega_4 \sim 673 \text{ cm}^{-1}$ C-H bend
- $\omega_2 \sim 993 \text{ cm}^{-1}$ C-C stretch

«hard»



«soft»

→ Collisional-vibrational excitation?

Benzene scattering: Key questions

- Translational energy:

→ TD vs. IS: Do the molecules equilibrate with the surface?

- Rotational excitation:

→ Is there transfer from translational to rotational energy?

- Vibrational excitation:

→ Can we observe collisional-vibrational excitation?

Benzene scattering: Key questions

- Translational energy:

→ TD vs. IS: Do the molecules equilibrate with the surface?

- Rotational excitation:

→ Is there transfer from translational to rotational energy?


- Vibrational excitation:

→ Can we observe collisional-vibrational excitation?

Dynamics of inelastic collision


tunable collision energy:
 $\langle E_{\text{inc}} \rangle \sim 8 \text{ kcal mol}^{-1} \sim 2800 \text{ cm}^{-1}$

Vibronic transitions in benzene (D_{6h})



1 + 1 REMPI scheme to ionize
vibrational ground state of
benzene

Vibronic transitions in benzene (D_{6h})




Investigate excited vibrational states of ground state benzene after the scattering process

Y: normal mode

- [2] G. Herzberg, *Mol. Spectra & Mol. Structure, Volume III*, Krieger, 1991
- [3] E. Riedle et al., *JCP*, **91**, 4555, 1989
- [4] M. Okruss et al., *JMS*, **193**, 293, 1999

Vibronic transitions in benzene (D_{6h})



Quantitative comparison
of the population
in each vibrational mode:


Line strength:

$$\begin{aligned} & (\langle ev(S_1 \ v_6 \ ^1=1 \ v_Y \ ^1=1) | \mu | ev(S_0 \ v_Y=1) \rangle)^2 \\ & \sim (\langle ev(S_1 \ v_6 \ ^1=1) | \mu | ev(S_0) \rangle \\ & \quad \times \langle v_Y \ ^1=1 | v_Y=1 \rangle)^2 \end{aligned}$$

both electronic states
very similar in structure
→ $\langle v_Y \ ^1=1 | v_Y=1 \rangle$
very similar for all Y

Translational energy distribution


Velocity-map image of vibrational ground state of benzene:



- heated Au surface (500 K)
- $S_1 \leftarrow S_0 6_0^1$ transition
- 2% benzene in 70% Ne/30% He
- incident beam ~ 8 kcal/mol
- TD and IS overlap strongly

Translational energy distribution


→ Integrated velocity-map image:



- $T \sim 1000 \text{ K}$
- significantly hotter than surface
- IS component dominant
- almost no forward-scattering
- ~20% of translational energy retained

Rotational excitation of ground state


Integrate over translational distribution and scan detection frequency



- rotational distribution hotter than surface
- not a single rotational T
- not equilibrated with surface
- dynamics during scattering process
- 30% of incident beam energy in rotation
- energy transfer with surface

Vibrational excitation in the scattering

REMPI spectrum of the integrated scattered flux:



- significant vibrational excitation
- not all low-energy modes equally excited

Conclusions & Outlook

- benzene is a ‹soft› molecule:
 - almost no forward-scattering of scattered molecules
 - impulsively-scattered component is dominant, no equilibrium
 - efficient transfer from translational to rotational energy
 - collisional-vibrational energy transfer, complex dynamics
- next steps:
 - better understanding of rotational and vibrational excitation
 - increase collision energy

Thank you!

20	E2u	CCC bend perp	405
19	E2u	C-H bend perp	970
18	E2g	CCC bend par	608.13
17	E2g	C-H bend par	1177.78
16	E2g	C-C stretch	1591.33
15	E2g	C-H stretch	3056.7
14	E1u	C-H bend par	1038.27
13	E1u	C-C stretch	1483.98
12	E1u	C-H stretch	3047.91
11	E1g	C-H bend perp	845
10	B2u	C-H bend par	1149.7
9	B2u	C-C stretch	1309.4
8	B2g	CCC bend perp	703
7	B2g	C-H bend perp	992
6	B1u	CCC bend par	1010
5	B1u	C-H stretch	3059
4	A2u	C-H bend perp	673
3	A2g	C-H bend par	1366.6
2	A1g	C-C stretch	993.06
1	A1g	C-H stretch	3073.94