

Ozone Photolysis: strong Isotopologue selectivity in the stratosphere.

F. Gatti⁽¹⁾, S. Ndengué⁽³⁾, H.-D. Meyer⁽²⁾, R. Jost⁽³⁾.

(1) CTMM Institut Charles Gerhardt UMR-CNRS 5253

University of Montpellier, France

(2) Institute for Physical Chemistry

University of Heidelberg, Germany


(3) Laboratoire de Spectrométrie Physique

University of Grenoble, France


Unusual isotopic constitution of Ozone

- Ozone has an unusual ^{17}O and ^{18}O isotopic constitution in the atmosphere
- Due to ozone formation process but also photolysis especially in the stratosphere (estimated previously at 25 % of the isotopic enrichment)
- Studied previously
M. C. Liang, F. W. Irion, Weibel, J. D. Blake, C. E. Miller and Y. L. Yung, J. Geophys. Res. **2006**, *111* D02302.
D. Krankowsky and K. Lämmerzahl and K. Mauersberger and J. Janssen and B. Tuzson and T. Röckmann, J. Geophys. Res. **2007**, *112* D08301.
- New calculations using Potential Energy Surfaces (PESs) and Dipole Moment Surfaces by Schinke and co-workers and the Heidelberg Multi-Configuration Time-Dependent Hartree (MCTDH) package
<http://www.pci.uni-heidelberg.de/cms/mctdh.html>

Focus on the Huggins band



Calculation of the Hartley band



S. Ndengué and F. Gatti and R. Schinke and H.-D. Meyer and R. Jost, JPCA (2010) 114 9855.

Five isotopologues



Symmetric $^{16}\text{O}^{17}\text{O}^{16}\text{O}$ (676), $^{16}\text{O}^{18}\text{O}^{16}\text{O}$ (686)


Asymmetric $^{16}\text{O}_2^{17}\text{O}$ (667), $^{16}\text{O}_2^{18}\text{O}$ (668)

668 can dissociate into either 66 + 8 or 68 + 6


The dissociated Oxygen atom can be triplet or singlet

We neglect the dissociation into three atoms of oxygen


PES of the B $3^2A'$ state



Calculation of the vibrational B states




States (2,3,0) and (2,0,1)



States 95 and 100


Absorbed cross sections (Huggins)



Experimental cross section at 218 K : J. Brion (Reims)


Theoretical values energy shift (to lower energy) of 450 cm^{-1} and a 285/234 factor of rescaling.

Cross section ratios




S. Ndengué and R. Schinke and F. Gatti H.-D. Meyer and R. Jost, JPCA (2012) 116
12260.

Asymmetric isotopologues



Singlet-Triplet Oxygen atom



S. Ndengué and R. Schinke and F. Gatti H.-D. Meyer and R. Jost, JPCA (2012) 116
12271.

Isotopic enrichment

Photolysis rate coefficient:

$$(1) \quad J_{abc} = \int X S_{abc}(E) Ac.Flux(E, h) dE$$


Tropospheric Ultraviolet-Visible model version 5:

Sasha Madronich and G. Weller, TUV program J. Atmos. Chem. (1990) 10 289
averaged over latitudes and seasons.

The enrichment of an abc isotopologue :

$$(2) \quad I_{abc} = -100 * (J_{abc}/J_{666} - 1)$$

Isotopic enrichment



Comparison with

M. C. Liang, F. W. Irion, Weibel, J. D. Blake, C. E. Miller and Y. L. Yung, J. Geophys. Res. 2006, 111 D02302.

Conclusion

The contribution of the photolysis process to the ozone and atomic oxygen isotope enrichment should be increased respectively by 25 % and 50 % for the asymmetric and symmetric isotopologues

Necessity to combine these results with the formation process to have the global enrichment

Acknowledgements:

- **Grenoble: S. Ndengué and R. Jost**
- **Heidelberg: H.-D. Meyer**
- **Göttingen: R. Schinke**
- Funding : ANR IDEO (NT09_436466), DFG