


# Detection and Formation of Interstellar c-C<sub>3</sub>D<sub>2</sub>

Silvia Spezzano

S. Schlemmer, S. Brünken and P. Schilke (Uni Köln), K. M. Menten (MPIfR, Bonn),  
M. C. McCarthy (Harvard), P. Caselli (Leeds), Y. Aikawa (Kobe), L. Bizzocchi  
(Lisboa), S. Trevino (IRAM)

**68<sup>th</sup> International Symposium on Molecular Spectroscopy**




Max-Planck-Institut  
für Radioastronomie

**IMPRS**  
astronomy &  
astrophysics  
Bonn and Cologne

Universität zu Köln



# Introduction - *Deuterium* -





# Introduction - Gas vs Grains -

Multiply deuterated species :

**$\text{H}_3^+$**  ( $\text{H}_2\text{D}^+$ ,  $\text{D}_2\text{H}^+$ ,  $\text{D}_3^+$ )

**Methanol** ( $\text{CH}_2\text{DOH}$ ,  $\text{CHD}_2\text{OH}$ ,  $\text{CD}_3\text{OH}$ )

**Ammonia** ( $\text{NH}_2\text{D}$ ,  $\text{NHD}_2$ ,  $\text{ND}_3$ )


**Formaldehyde** ( $\text{HDCO}$ ,  $\text{D}_2\text{CO}$ )

**Hydrogen Sulfide** ( $\text{HDS}$ ,  $\text{D}_2\text{S}$ )

**Thioformaldehyde** ( $\text{HDCS}$ ,  $\text{D}_2\text{CS}$ )



# Introduction - $c\text{-C}_3\text{H}_2$ -



First identification in 1985 (Thaddeus et al.)

First detection of  $c\text{-C}_3\text{HD}$  in 1986 (Bell et al.)

**HIGH DEUTERATION :**


$c\text{-C}_3\text{HD}/c\text{-C}_3\text{H}_2 \cong 1/5$  in TMC-1 (Gerin et al., 1987)



Horsehead Nebula  
Credit: ESO



M82  
Credit: ESO




NGC 7027  
Credit: ESO





# What makes c-C<sub>3</sub>H<sub>2</sub> unique?


- \* Widespread
- \* Possibility of double deuteration
- \* Solely gas phase chemistry




Empirical equilibrium structure of c-C<sub>3</sub>H<sub>2</sub>



# Laboratory detection of c-C<sub>3</sub>D<sub>2</sub>



$1_{10} - 1_{01}$  transition near 17 GHz  
 Fourier Transform microwave spectrometer  
 (8 - 40 GHz)



Ortho-para doublet near 233 GHz  
 Double pass absorption spectrometer  
 (200 - 320 GHz)

*Spezzano et al., ApJ Supp. S. (2012)*



# Observations

36 h observing time, end of September

Sources: starless cores TMC-1C and L1544

EMIR receivers in E090 configuration

Lines observed:

c-C<sub>3</sub>H<sub>2</sub> @ 84 and 85 GHz

c-<sup>13</sup>C<sub>3</sub>H<sub>2</sub> (off axis) @ 84 GHz

c-C<sub>3</sub>HD @ 95 and 104 GHz

c-C<sub>3</sub>D<sub>2</sub> @ 94, 97 and 108 GHz

Frequency switching mode

FTS spectrometer with 50kHz resolution

(0.015 km s<sup>-1</sup> at 3mm)




IRAM 30 m telescope  
Pico Veleta (Spain)






# Results

L1544



TMC-1C



- All favourable transitions of  $c\text{-C}_3\text{D}_2$  available in the covered frequency range have been detected
- The rest frequencies employed have laboratory accuracy, and in both sources the line shapes and velocities are in agreement with each other





# Results

Table 1. Abundance ratios of deuterated molecules in TMC-1C and L1544

	TMC-1C	L1544
$[c\text{-C}_3\text{D}_2]/[c\text{-C}_3\text{H}_2]$	(0.4 - 0.8)%	(1.2 - 2.1)%
$[c\text{-C}_3\text{D}_2]/[c\text{-C}_3\text{HD}]$	(3 - 15)%	(7- 17)%
$[c\text{-C}_3\text{HD}]/[c\text{-C}_3\text{H}_2]$	(5 - 13)%	(12 - 17)%
$[\text{D}_2\text{CO}]/[\text{H}_2\text{CO}]$	-	4% <sup>a</sup>
$[\text{DCO}^+]/[\text{HCO}^+]$	2% <sup>b</sup>	4% <sup>c</sup>
$[\text{N}_2\text{D}^+]/[\text{N}_2\text{H}^+]$	8% <sup>b</sup>	20% <sup>c</sup>
$[\text{NH}_2\text{D}]/[\text{NH}_3]$	1% <sup>b</sup>	13% <sup>d</sup>

<sup>a</sup>Bacmann (2003), <sup>b</sup>Tiné (2000), <sup>c</sup>Caselli (2002), <sup>d</sup>Shah (2001)

*Spezzano et al., ApJL (2013)*



# Results

D/H ratio of C<sub>3</sub>H<sub>2</sub> in prestellar cores was calculated using the network model of Aikawa et al (2012)

Physical structure of the core: collapsing core model of Aikawa (2005) and also a static model of L1544 from Keto and Caselli (2010)

For CO depletion factors consistent with those of the 2 objects the calculated column density ratio of C<sub>3</sub>D<sub>2</sub>/C<sub>3</sub>H<sub>2</sub> is  $\sim 10^{-2}$

	TMC-1C	L1544
$[c\text{-C}_3\text{D}_2]/[c\text{-C}_3\text{H}_2]$	(0.4 - 0.8)%	(1.2 - 2.1)%


There is no need for any reaction on grains to account for the observed D/H ratio: the deuteration of cyclopropenylidene can be solely explained by gas phase reactions

*Spezzano et al., ApJL (2013)*





# Results



→ proton/deuteron transfer SLOW PROCESS

→ dissociative recombination with  $e^-$  FAST PROCESS



# Summary

- ♦ First interstellar detection of doubly deuterated cyclopropenylidene
- ♦  $c\text{-C}_3\text{D}_2$  ideal probe for the study of gas phase deuteration reactions
- ♦ High resolution spectroscopy was a prerequisite for the astronomical detection





# Summary

- ♦ First interstellar detection of doubly deuterated cyclopropenylidene
- ♦  $c\text{-C}_3\text{D}_2$  ideal probe for the study of gas phase deuteration reactions
- ♦ High resolution spectroscopy was a prerequisite for the astronomical detection

# Future

- ♦ Map
- ♦ Spectroscopy of isotopic species of  $\text{C}_3\text{H}_3^+$



# Acknowledgements

International Max Planck Research School for Astronomy and  
Astrophysics in Bonn and Cologne

SFB 956

Fondazione Angelo della Riccia Fellowship







**68<sup>th</sup> International Symposium on  
Molecular Spectroscopy**  
**June 18<sup>th</sup> 2013**

**Silvia Spezzano**  
**I. Physikalisches Institut**  
**Universität zu Köln**

**Universität zu Köln**



# Results

Column densities and optical depth have been calculated in the following way

$$\tau_0 = \ln \left( \frac{J(T_{ex}) - J(T_{bg})}{J(T_{ex}) - J(T_{bg}) - T_{mb}} \right)$$

$$J(T) = \frac{h\nu}{k} \left( e^{\frac{h\nu}{kT}} - 1 \right)^{-1}$$

$$N_{up}^{thin} = \frac{8\pi\nu^3 \sqrt{\pi} \Delta\nu \tau_0}{c^3 A_{ul} 2\sqrt{\ln 2} \left( e^{\frac{h\nu}{kT}} - 1 \right)}$$

$$N_{tot} = \frac{N_u Q_{rot}(T_{ex})}{g_u e^{\left(\frac{E_u}{kT_{ex}}\right)}}$$

$A_{ul}$  and  $B_{ul}$  : Einstein coefficients

$Q_{rot}(T)$  : partition function at given  $T$

$g_u$  : degeneracy of the upper level

$T_{ex}$  (TMC-1C) = 7 K

$T_{ex}$  (L1544) = 5 K

