



CALIFORNIA INSTITUTE OF TECHNOLOGY

AN EMPIRICAL APPROACH TO OBTAINING ACCURATE MOLECULAR ROTATIONAL CONSTANTS FOR ISOTOPICALLY- SUBSTITUTED SPECIES FROM AB INITIO CALCULATIONS

Brett A. McGuire • P. Brandon Carroll • Steven T. Shipman • Geoffrey A. Blake

68th International Symposium on Molecular Spectroscopy - June 19, 2013




INTRODUCTION - ISOTOPLOGUES

Why study isotopic substitution?

Zyankun, A. M., 2011, J Analytic. Chem., 66, 1243
Carroll, McGuire, Zaleski et al., 2013, J Mol Spec, 284-285, 21



Why study isotopic substitution?




Zyankun, A. M., 2011, J Analytic. Chem., 66, 1243
Carroll, McGuire, Zaleski et al., 2013, J Mol Spec, 284-285, 21





INTRODUCTION - ISOTOPOLOGUES

Why study isotopic substitution?



Zyankun, A. M., 2011, J Analytic. Chem., 66, 1243
Carroll, McGuire, Zaleski et al., 2013, J Mol Spec, 284-285, 21

Why study isotopic substitution?





Zyankun, A. M., 2011, J Analytic. Chem., 66, 1243
Carroll, McGuire, Zaleski et al., 2013, J Mol Spec, 284-285, 21




INTRODUCTION - ISOTOPOLOGUES

Why study isotopic substitution?



Zyankun, A. M., 2011, J Analytic. Chem., 66, 1243
Carroll, McGuire, Zaleski et al., 2013, J Mol Spec, 284-285, 21

More accurate rotational constants result in better initial predictions of rotational transition frequencies






INTRODUCTION - ISOTOPOLOGUES

More accurate rotational constants result in better initial predictions of rotational transition frequencies

WH16

6:00
1000 MP





What's wrong with what we have now?

- Basic computational methods return
equilibrium (R_e) geometry



What's wrong with what we have now?

- Basic computational methods return equilibrium (R_e) geometry
- R_0 geometry → much more expensive



What's wrong with what we have now?

- Basic computational methods return equilibrium (R_e) geometry
- R_0 geometry → much more expensive
- Can do extremely well... for a price



INTRODUCTION - GOAL

Most accurate isotopologue structure with least computational expense (effort)



INTRODUCTION - GOAL

Most accurate isotopologue structure with least computational expense (effort)

- Parent species data → “easily” obtained



INTRODUCTION - GOAL

Most accurate isotopologue structure with least computational expense (effort)

- Parent species data → “easily” obtained
- Perform “simple” optimizations



INTRODUCTION - GOAL

Most accurate isotopologue structure with least computational expense (effort)

- Parent species data → “easily” obtained
- Perform “simple” optimizations
- Combine to quickly obtain useful predictions



METHODS - THE METHOD

I. Optimize structure at given Level of Theory/Basis Set

- Calculate parent species rotational constants



METHODS - THE METHOD

- I. Optimize structure at given Level of Theory/Basis Set
 - Calculate parent species rotational constants
2. Assume no geometry change upon substitution
 - Calculate isotopologue rotational constants



METHODS - THE METHOD

1. Optimize structure at given Level of Theory/Basis Set
 - Calculate parent species rotational constants
2. Assume no geometry change upon substitution
 - Calculate isotopologue rotational constants
3. Scale isotopologue rotational constants



METHODS - THE METHOD

$$B_{\text{scaled}} = \frac{B_{\text{exp}} \text{ (parent)}}{B_{\text{calc}} \text{ (parent)}} \times B_{\text{calc}}$$



B U T W A I T ...

Hold on, doesn't everyone do this already?



B U T W A I T ...

Hold on, doesn't everyone do this already?

Probably?



Journal of Molecular Spectroscopy 284-285 (2013) 21–28


The pure rotational spectrum of glycolaldehyde isotopologues observed in natural abundance

P. Brandon Carroll^a, Brett A. McGuire^a, Daniel P. Zaleski^b, Justin L. Neill^b, Brooks H. Pate^{b,*}, Susanna L. Widicus Weaver^{a,*}

New Techniques in Microwave Spectroscopy

Steven T. Shipman and Brooks H. Pate

Department of Chemistry, University of Virginia, Charlottesville, VA, USA





Levels of theory

MP2 • B3LYP • M062X • CCSD

Basis sets

6-31+G • 6-311++G(d,p) • cc-pVDZ • aug-cc-pVTZ



METHODS - MOLECULES


2




3




4




6





7



8




9






RESULTS - OVERALL






RESULTS - OVERALL





RESULTS - ANHARMONIC COMPARISON

Comparision of Effects of Including Anharmonic Ro-Vibrational Interactions





RESULTS - HCl

H³⁷Cl - Dunham

Lit = 312 519.12 MHz

$$B(\text{H}^{37}\text{Cl}) = \frac{\mu(\text{H}^{35}\text{Cl})}{\mu(\text{H}^{37}\text{Cl})} \times B(\text{H}^{35}\text{Cl}) = 312\ 519.35\ \text{MHz}$$

(0.23 MHz)



RESULTS - HCl

H³⁷Cl - Dunham

Lit = 312 519.12 MHz

$$B(\text{H}^{37}\text{Cl}) = \frac{\mu(\text{H}^{35}\text{Cl})}{\mu(\text{H}^{37}\text{Cl})} \times B(\text{H}^{35}\text{Cl}) = 312\ 519.35\ \text{MHz}$$

(0.23 MHz)

H³⁷Cl - This work

$$B(\text{H}^{37}\text{Cl}) = 312\ 518.81\ \text{MHz}$$

(0.31 MHz)



RESULTS - HCl

H³⁷Cl - Dunham

Lit = 312 519.12 MHz

$$B(\text{H}^{37}\text{Cl}) = \frac{\mu(\text{H}^{35}\text{Cl})}{\mu(\text{H}^{37}\text{Cl})} \times B(\text{H}^{35}\text{Cl}) = 312\ 519.35\ \text{MHz}$$

(0.23 MHz)

H³⁷Cl - This work

$$B(\text{H}^{37}\text{Cl}) = 312\ 518.81\ \text{MHz}$$


(0.31 MHz)

Only this accurate at
CCSD



RESULTS - TRIATOMICS


Error in Predictions for Triatomics






RESULTS - TRIATOMICS

Error in Predictions for Triatomics





RESULTS - TRIATOMICS





RESULTS - OVERALL

Takeaway key points

- **Be cautious with deuterium**
-



RESULTS - OVERALL

Takeaway key points

- **Be cautious with deuterium**
- Best consistent improvement
CCSD/aug-cc-pVTZ



RESULTS - OVERALL

Takeaway key points

- **Be cautious with deuterium**
- Best consistent improvement
CCSD/aug-cc-pVTZ
- Most time efficient improvement
MP2/6-311++G(d,p)



RESULTS - OVERALL

Takeaway key points

- **Be cautious with deuterium**
- Best consistent improvement
CCSD/aug-cc-pVTZ
- Most time efficient improvement
B3LYP/6-31+G



EXTENSIONS - AUTOMATED ANALYSIS


Assigning $CH_3C(O)OD$



Assigning $CH_3C(O)OD$

Start with $|_{1,0} - |_{0,1}$ transition

Scale by abundance: Intensity = 0.057






Assigning $CH_3C(O)OD$

Start with $|_{1,0} - |_{0,1}$ transition

Scale by abundance: Intensity = 0.057

Lowest *B*-type: Freq = *A* - *C*

Error = Error (*A*) + Error (*C*)



Assigning $CH_3C(O)OD$

Start with $|_{1,0} - |_{0,1}$ transition

Scale by abundance: Intensity = 0.057

Lowest *B*-type: Freq = *A* - *C*

Error = Error (*A*) + Error (*C*)


Error (Unratioed)

165 MHz

Error (Ratioed)


6 MHz

B3LYP/aug-cc-pVQZ






EXTENSIONS - AUTOMATED ANALYSIS






EXTENSIONS - AUTOMATED ANALYSIS






EXTENSIONS - AUTOMATED ANALYSIS






EXTENSIONS - AUTOMATED ANALYSIS






EXTENSIONS - AUTOMATED ANALYSIS





EXTENSIONS - AUTOMATED ANALYSIS





ACKNOWLEDGMENTS

Special Thanks

Joel Bowman

Susanna Widicus Weaver

Thomas Miller

Funding



Take Home Message

We have presented a computationally inexpensive method for obtaining accurate rotational constants using mixture of experiment and theory



ACKNOWLEDGMENTS

Special Thanks

Joel Bowman

Susanna Widicus Weaver

Thomas Miller

Funding



Take Home Message


We have presented a computationally inexpensive method for obtaining accurate rotational constants using mixture of experiment and theory



EXTENSIONS - VIBRATIONAL MODES

What happens if we apply the same methodology to calculated vibrational frequencies of $\text{H}_2\text{CO}/\text{D}_2\text{CO}$?

Error in Fundamental Vibrational Predictions for D_2CO



Requires identical symmetry!