

CARS STUDY OF DEUTERIUM CLUSTERS STABILIZED IN SOLID HELIUM

EUGENE B. GORDON, *Institute of Problems of Chemical Physics, Chernogolovka, 142432 Moscow Region, Russia* ; TAKAYUKI KUMADA, MASAZUMI ISHIGURO, and YASUYUKI ARATONO , *Advanced Science Research Center, Japan Atomic Energy Research Institute, Tokai, Ibaraki 319-1195, Japan*.

By using the technique recently proposed^a deuterium clusters have been firstly introduced into the matrix being more inert than hydrogen itself^b. The known effect of Q₁(1) line dramatic predominance over Q₁(0) one turned out to be even more expressed in CARS than in common Raman scattering (RS) - being equal to 1 in a gas the ratio of probabilities for nearly pure o-D₂ crystal was found to be as high as 10,000 in stimulated RS whereas in common RS it has been known to be only 50. We have shown that the effect appears just as the phonon band is to be formed in crystallite, i.e. for the clusters contained more than 10³ molecules. The coherent interaction of D₂ resonant line with non-resonant CARS signal from solid helium has been observed.

^aE. B. Gordon, A. Usenko and G. Frossati *J. Low Temp. Phys.* **130**(1-2), 15, 2003

^bE. B. Gordon, G. Frossati, A. Usenko, Y. Aratono and T. Kumada *Physica B* (in press)