

## FOURIER TRANSFORM MICROWAVE AND INFRARED SPECTROSCOPIC INVESTIGATION OF PROPIOLACTONE

ZIQU CHEN and JENNIFER VAN WIJNGAARDEN, *Department of Chemistry, University of Manitoba, Winnipeg MB R3T 2N2 Canada.*

The pure rotational spectrum of the four-membered ester ring propiolactone ( $C_3H_4O_2$ ) has been measured in a supersonic jet between 7 and 22 GHz using Fourier transform microwave (FTMW) spectroscopy. For the normal isotopologue, a total of 19 *a*- and *b*-type transitions have been recorded. Fifteen transitions due to three different  $^{13}C$  isotopologues have also been observed. The microwave spectrum was analyzed to obtain an improved set of ground state rotational constants in comparison to earlier microwave experiments<sup>a</sup>. The new set of rotational parameters was used to predict the rovibrational band structure of the lowest frequency modes of propiolactone. A total of 12 vibrational band origins have been observed between 400 and 1500  $cm^{-1}$  using the far infrared beamline of the Canadian Light Source coupled to a Bruker IFS125HR spectrometer. The spectra were recorded with a resolution of 0.000969  $cm^{-1}$  and although the intensities of the bands vary, 9 bands are of sufficient quality for complete rovibrational assignment. The progress of the assignment of this rich spectrum will be discussed.

---

<sup>a</sup>D. W. Boone, C O. Britt and J. E. Boggs *J. Chem. Phys.* **43** (1190), 1965.