

ISOTOPOLOGUE-SENSITIVE DETECTION USING CHIRPED-PULSE FT-MW SPECTROSCOPY: MINOR SPECIES OF PROPOFOL

ALBERTO LESARRI, *Departamento de Química Física y Química Inorgánica, Facultad de Ciencias, Universidad de Valladolid, E-47011 Valladolid, Spain*; JUSTIN NEILL, MATT MUCKLE, STEVEN T. SHIPMAN, BROOKS H. PATE and RICHARD D. SUENRAM, *Department of Chemistry, University of Virginia, Charlottesville, VA 22904, USA*; WALTHER CAMINATI, *Dipartimento di Chimica "G. Ciamician", Università di Bologna, I-40126 Bologna, Italy*.

The capabilities of chirped-pulse FT-microwave spectroscopy to achieve full-bandwidth (11 GHz) isotopologue-sensitive detection have been tested on the 13-heavy atoms molecule of propofol (2,6-diisopropylphenol). The analysis of the rotational spectrum using moderate signal averaging (10 k FIDs) had previously detected the presence of two conformers arising from the combined internal rotations of the hydroxyl and the two isopropyl groups^a. In the new experiment reported here 600 k FIDs were coherently averaged, using three pulsed nozzle sources and reading multiple FIDs per sample injection cycle to reduce the total acquisition time and sample consumption. The new spectrum revealed a very large number of weak transitions, suggesting that full-band ¹³C sensitivity had been surpassed. The new data have resulted in the assignment of a third conformer of propofol, followed by all twelve ¹³C-monosubstituted species in natural abundance for the most stable conformer. The isotopic information confirmed the molecular structure for the preferred conformation of propofol, validating the *ab initio* predictions for this compound. The potential function for the OH internal rotation has been determined using a flexible model.

^aA. Lesarri, S. T. Shipman, G. G. Brown, L. Alvarez-Valtierra, R. D. Suenram and B. H. Pate, 63rd OSU Int. Symp. On Mol. Spectrosc., Columbus, OH, 2008, RH07