

A COMPUTATIONAL INVESTIGATION OF $c\text{-C}_3\text{H}_2\dots\text{HX}$ ($X = \text{F}, \text{Cl}, \text{Br}$) H-BONDED COMPLEXES

PRADEEP R. VARADWAJ, ARPITA VARADWAJ, GILLES H. PESLHERBE, *Centre for Research in Molecular Modeling & Department of Chemistry and Biochemistry, Concordia University, Montreal, QC, Canada.*

Cyclopropenylidene ($c\text{-C}_3\text{H}_2$) is of significant importance in interstellar chemistry and synthetic chemistry (e.g., transition metal and organic catalysis). Because of its peculiar structure, $c\text{-C}_3\text{H}_2$ can act as a hydrogen-bond donor or acceptor. In order to gain insight into this feature, the ground-state potential energy surfaces of singlet $c\text{-C}_3\text{H}_2$ complexed with hydrogen halides HX ($X = \text{F}, \text{Cl}, \text{Br}$) have been explored extensively by density-functional theory (B3LYP) and *ab initio* quantum chemistry (MP2) with a variety of basis sets, cc-pVxZ and aug-cc-pVxZ ($x = \text{D}, \text{T}$). The complexes characterized have the carbenic end of $c\text{-C}_3\text{H}_2$ H-bonded to HX, with some proton transfer occurring, the extent of which follows the order $\text{HF} < \text{HCl} < \text{HBr}$. Accompanying the complex formation are the dipole moment enhancement, the charge transfer, red shifts of the HX vibrational stretching frequencies together with the significant enhancement of band intensity and concomitant HX bond elongation. The nature of H-bonding in these complexes has been explored, based on energy decomposition schemes and the Baders quantum theory of atoms-in-molecules, with the conclusion that $c\text{-C}_3\text{H}_2$ is a strong H-bond acceptor with respect to the hydrogen halides.