

ROTATIONAL AND HYPERFINE ANALYSIS OF THE $a^1\Delta_g \leftarrow X^3\Sigma_g^-$ BAND OF ^{17}O -CONTAINING ISOTOPOLOGUES OF OXYGEN MEASURED BY CRDS AT ROOM AND LIQUID NITROGEN TEMPERATURES

O. M. LESHCHISHINA, S. KASSI, *Université de Grenoble, CNRS UMR 5588, LIPHY, 38041 Grenoble, France*; I. E. GORDON, *Harvard-Smithsonian Center for Astrophysics, Atomic and Molecular Physics Division, Cambridge MA 02138-1516, USA*; S. YU, *Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109, USA*; A. CAMPARGUE, *Université de Grenoble, CNRS UMR 5588, LIPHY, 38041 Grenoble, France*.

The $a^1\Delta_g \leftarrow X^3\Sigma_g^-$ system of the $^{16}\text{O}^{17}\text{O}$, $^{17}\text{O}^{18}\text{O}$ and $^{17}\text{O}_2$ isotopologues of oxygen was studied by high sensitivity CW-Cavity Ring Down Spectroscopy. The spectra of a ^{17}O highly enriched sample were recorded at room temperature between 7640 and 7917 cm^{-1} and at liquid nitrogen temperature in the 7876-7893 cm^{-1} region. The magnetic dipole (0-0) band was observed for all three ^{17}O -containing isotopologues. At liquid nitrogen temperature some of the transitions were observed with partially resolved hyperfine splitting due to the ^{17}O nuclear spin, allowing determination of the hyperfine constants. The electric quadrupole (0-0) band and the (1-1) magnetic dipole hot band were also observed for the $^{16}\text{O}^{17}\text{O}$ and $^{17}\text{O}_2$ species. The rotational and hyperfine spectroscopic parameters of the $X^3\Sigma_g^-$ and $a^1\Delta_g$ states of the three studied isotopologues were derived from global fit of the measured line positions and microwave and Raman measurements available in the literature. The rotational constants of the $a^1\Delta_g$ ($v=0, 1$) states of $^{17}\text{O}_2$ are determined for the first time.