

ARE AB INITIO QUANTUM CHEMISTRY METHODS ABLE TO PREDICT VIBRATIONAL STATES UP TO THE DISSOCIATION LIMIT FOR MULTI-ELECTRON MOLECULES CLOSE TO SPECTROSCOPIC ACCURACY?

PÉTER G. SZALAY, *Eötvös Loránd University, Budapest, Hungary*; FILIP HOLKA, *Slovak University of Technology, Trnava, Slovak Republic*; JULIEN FREMONT, MICHAEL REY, VLADIMIR G. TYUTEREV, *Reims University, Reims, France*.

The aim of the study was to explore the limits of *ab initio* methods towards the description of excited vibrational levels up to the dissociation limit for molecules having more than two electrons. To this end a high level *ab initio* potential energy function was constructed for the four-electron LiH molecule in order to accurately predict a complete set of bound vibrational levels corresponding to the electronic ground state. It was composed from: a) an *ab initio* non-relativistic potential obtained at the MR-CISD level including size-extensivity corrections and quintuple-sextuple  $\zeta$  extrapolation of the basis, b) MVD (Mass-velocity-Darwin) relativistic corrections obtained at icMR-CISD/cc-pwCV5Z level, and c) DBOC (Diagonal Born-Oppenheimer correction) obtained at the MR-CISD/cc-pwCVTZ level. Finally, the importance of non-adiabatic effects was also tested by using atomic masses in the vibrational kinetic energy operator and by calculation of non-adiabatic coupling by *ab initio* methods.

The calculated vibrational levels were compared with those obtained from experimental data [J.A. Coxon and C.S. Dickinson, *J. Chem. Phys.*, 2004, **121**, 9378]. Our best estimate of the potential curve results in vibrational energies with a RMS deviation of only  $\sim 1$   $\text{cm}^{-1}$  for the entire set of all empirically determined vibrational levels known so far. These results represent a drastic improvement over previous theoretical predictions of vibrational levels of  ${}^7\text{LiH}$  up to dissociation,  $D_0$ , which was predicted to be  $19594$   $\text{cm}^{-1}$ .

In addition, rotational levels have also been calculated. The RMS deviation between our *ab initio* calculations and empirical results by Coxon and Dickinson for rotational spacings  $\Delta E = E(v, J = 1) - E(v, J = 0)$  over all available vibrational states of  ${}^7\text{LiH}$  from  $v = 0$  to  $v = 20$  is  $0.010$   $\text{cm}^{-1}$  (with nuclear masses) and  $0.006$   $\text{cm}^{-1}$  (with atomic masses). Note that for high vibrational states with  $v > 6$  this falls within the uncertainty of the measurements.