

THE MILLIMETERWAVE SPECTRUM OF *n*-BUTYL CYANIDE

MATTHIAS H. ORDU, HOLGER S. P. MÜLLER, FRANK LEWEN, STEPHAN SCHLEMMER,
I. Physikalisches Institut, Universität zu Köln, Zùlpicher Str. 77, 50937 Köln, Germany; MARC NUÑEZ,
and ADAM WALTERS, *IRAP: Université de Toulouse, UPS-OMP, CNRS; 9 Av. colonel Roche, BP 44346,*
31028 Toulouse cedex 4, France.

The rotational spectrum of *n*-butyl cyanide (C_4H_9CN) was measured between 75 and 130 GHz using a novel all-solid-state spectrometer with a total absorption path of 44 m. In the course of the analysis of the spectrum, about 3000 transitions were assigned and a full set of quartic centrifugal distortion parameters with some sextic and octic terms could be determined for each of the three known conformers (anti-anti, anti-gauche(methyl end) and gauche(CN end)-anti).

The work was motivated by the fact that *n*-butyl cyanide is likely to be found in interstellar hot core environments. This is indicated by the discovery of *n*-propyl cyanide (C_3H_7CN), the next smaller alkyl cyanide, in the ISM^a. The increased accuracy of the model, which will be additionally extended by future laboratory measurements around 200 GHz, may now be employed for a prediction of the spectrum up to 300 GHz with a feasible uncertainty for astronomic line surveys.

Furthermore, there are two less abundant conformers, *cis*-gauche-gauche and *trans*-gauche-gauche, which have not yet been detected in the rotational spectrum^b. Due to the increased sensitivity of the new spectrometer, it seems possible now for the first time to identify their spectroscopic fingerprints in the recorded data.

^aA. Belloche, R. T. Garrod, H. S. P. Müller, K. M. Menten, C. Comito, and P. Schilke, *Astronomy & Astrophysics* **499**, 215 (2009)

^bR. K. Bohn, J. L. Pardus, J. August, T. Brupbacher, W. Jäger, *J. Mol. Struct.* **413–414**, 293 (1997)