

## VIBRATIONALLY HOT HCN IN THE LABORATORY AND IRC+10216

JOHN C. PEARSON<sup>a</sup>, SHANSHAN YU, HARSHAL GUPTA and BRIAN J. DROUIN, *Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Dr., Pasadena, CA 91109.*

HCN has historically been used as a tracer of the dense gas in the interstellar medium. The envelopes of carbon rich asymptotic giant branch stars are generally rich in HCN; however, the large and generally variable infrared flux emitted by the star enormously complicates the interpretation. HCN in IRC+10216 shows an enormous number of masers and lasers pumped by the central star and often enhanced by line overlaps with other abundant molecules such as acetylene in the infrared. A total of seven laser transitions including two previously unreported transitions associated with the 040 – 011 interacting bands have been observed. To understand the astronomical observations a study of the radio frequency discharge plasma of CH<sub>4</sub> and N<sub>2</sub> was performed. Rotational transitions of HCN in vibrational states up to 15,000 cm<sup>-1</sup> have been observed including inverted levels and a number of previously undetected states. The spectra from IRC+10216 and the laboratory are presented.

---

<sup>a</sup>A part of this work was performed at the Jet Propulsion Laboratory, California Institute of Technology under contract with the National Aeronautics and Space Administration. Copyright 2010© California Institute of Technology. All rights reserved.