

BREAKING THE SYMMETRY IN JAHN-TELLER ACTIVE MOLECULES BY ASYMMETRIC ISOTOPIC SUBSTITUTION: SPLITTING THE ZERO-POINT VIBRONIC LEVEL.

DMITRY G. MELNIK, JINJUN LIU, TERRY A. MILLER, *Laser Spectroscopy Facility, Department of Chemistry, The Ohio State University, 120 W. 18th Avenue, Columbus, Ohio 43210*; ROBERT F. CURL, *Department of Chemistry and Rice Quantum Institute, Rice University, Houston, Texas 77005*.

Analysis and understanding of the vibronic spectra of Jahn-Teller active molecules have been challenging tasks due to the complex interaction patterns between vibronic levels even if the density of vibronic states is not too high. An example of a spectral feature that draws attention is the splitting of the ground vibronic level, as reported for degenerate electronic states of asymmetrically deuterated C_6H_6 , C_5H_5 , CH_4^+ , and most recently, the methoxy isotopologues, CH_2DO and CHD_2O . Understanding such splittings plays a crucial role for the analysis of the vibronic spectra of these species. We have developed a simplified model emulating vibronic interactions involving a single doubly-degenerate vibrational mode and one totally symmetric mode in a degenerate electronic state. The extension of this model to more complex, realistic cases and comparison with experiment and quantum chemical calculations will also be presented.