

**Problem Set 4**  
**Due Tuesday Feb 14, 2012**

1) The decay of an unstable particle is described by the following probability density function in terms of the decay time ( $t$ ) and the particle's lifetime ( $\lambda$ ).

$$p(t, \lambda) = \frac{e^{-\frac{t}{\lambda}}}{\lambda}$$

Three measurements of  $t$  ( $t_1 = 7$  sec,  $t_2 = 3$  sec,  $t_3 = 4$  sec) are made.

- Write down the likelihood function for this problem.
  - Use the Maximum Likelihood Method to calculate the value of  $\lambda$  for this data set.
- 2) A theory states that the angular distribution of electrons from the decay of an unstable particle should have a probability distribution function of the form (both  $N$  and  $\alpha$  are constants):

$$p(\cos \theta) = N(1 + \alpha \cos^2 \theta)$$

An experiment measures ten examples of the decay of this unstable particle and finds the following values of  $\cos \theta$ : (-0.05, -0.15, -0.25, -0.35, 0.45, 0.55, 0.65, 0.75, 0.85, 0.95). For this problem the limits on  $\cos \theta$  are [-1, 1]. We wish to determine the value of  $\alpha$  using the Maximum Likelihood Method.

a) Use the normalization condition for a probability distribution function to show that:

$$N = \frac{1}{2(1 + \alpha / 3)}$$

- Write down the Likelihood Function for this problem.
- Make a plot of the Likelihood Function vs.  $\alpha$  for  $-1.5 < \alpha < 1.5$ . Use this plot to find the value of  $\alpha$  that maximizes the Likelihood Function.

3) We wish to determine the acceleration due to gravity ( $g$ ) using the following data and  $\mathbf{h} = 0.5gt^2$ .

a) Use the least squares technique to find the best value of  $g$ . Assume the error in each  $h$  (height) measurement is 0.01 m and the time is measured exactly. (See Taylor Problem 8.5)

$h$ (m)	$t$ (s)
0.05	0.1
0.44	0.3
1.23	0.5
2.40	0.7

- What is the value of the chi-square ( $\chi^2$ ) for this problem?
- How many degrees of freedom are there in this problem? (See Taylor Problem 12.14, part b))
- Estimate the probability to get a  $\chi^2$  per degree of freedom  $\geq$  what you obtain using parts b) and c).

4) Taylor P8.4, page 200.

5) Taylor P8.10, page 201. Just do the first part of the problem (weighted LSQ estimate of A and B). Skip everything after "Compare..."