

**Physics 416
Problem Set 2**

Due Thursday, October 28, 2004

- 1) Taylor, Problem 10.10, page 242. In addition answer the following:
 - d) On average how many patients survive?
 - e) The probability that all four patients survive.
- 2) Assuming a Gaussian probability distribution answer the following questions
(Use Tables in *Taylor Appendix A and/or B*):
 - a) What is the probability of a value lying more than 1.5σ from the mean?
 - b) What is the probability of a value lying $\geq 1.5\sigma$ above the mean?
 - c) What is the probability of a value lying $\leq 1.5\sigma$ below the mean?
 - d) What is the probability of a value, y , lying in the range $\mu - \sigma \leq y \leq \mu + 2\sigma$?
 - e) What is the probability of a value, y , lying in the range $\mu + \sigma \leq y \leq \mu + 2\sigma$?

For this problem μ is the mean of the Gaussian and σ is its standard deviation.

- 3) Taylor, Problem 5.12, page 156.
- 4) The sun emits an enormous number of neutrinos. Assume that 10^6 solar neutrinos uniformly pass through a square with an area of 1 m^2 each μsec . Inside the square is a neutrino detector with an area of 1 mm^2 . Assume Poisson statistics for this problem.
 - a) What is the average number of neutrinos going through the detector each μsec ?
 - b) What is the probability that no neutrinos go through the detector in a μsec ?
 - c) What is the probability that \geq two neutrinos go through the particle detector in a μsec ?
 - d) How big should the detector be (in mm^2) if we want ≥ 1 particles per μsec to pass through the detector with a probability of 99%?
- 5) Taylor, Problem 11.18, page 259.
- 6) Suppose a missile defense system destroys an incoming missile 95% of the time.
 - a) If an evil country launches 20 missiles what is the probability that the missile defense system will destroy all of the incoming missiles?
 - b) How many missiles have to be launched to have a 50% chance of at least one missile making it through the defense system?

Note: this problem can be done using either binomial or Poisson statistics.

- 7) According to quantum mechanics, the position (x) of a particle in a one dimensional box with dimensions $-L/2 \leq x \leq L/2$ (L constant) can be described by the following probability distribution function $p(x)$:

$$p(x) = A \cos^2[\pi x/L] \text{ for } -L/2 \leq x \leq L/2, \text{ and } 0 \text{ for all other } x.$$

- a) Find the normalization constant A in terms of L .
- b) Find the mean, mode, and median position of the particle in the box.
- c) Show that the variance (σ^2) of x is given by:

$$\sigma^2 = \left(\frac{L}{\pi}\right)^2 \frac{\pi^2 - 6}{12}$$

d) What is the probability of finding the particle in the region: $L/4 \leq x \leq L/2$?

8) Suppose 100 six sided dice are tossed. Assume that the faces are labeled by one through six dots. Let Y_i be the number of dots on the i th ($i=1$ to 100) die.

a) What is the average number of dots expected for a single dice?

b) What is the variance of the numbers of dots expected for a single dice?

c) Use the Central Limit Theorem to estimate the probability that the sum of the Y_i 's exceeds 400.

9) In the Bohr theory of the structure of the hydrogen atom the energies of the various quantum states are given by:

$$E_n = - \frac{me^4}{2N^2\hbar^2}$$

With: m the mass of the electron

e the electric charge of the electron

\hbar Planck's constant divided by 2π

If: $\sigma_m/m = 0.1\%$ (i.e. the mass is known to 0.1%)

$\sigma_e/e = 0.2\%$ (i.e. the charge is known to 0.2%)

$\sigma_{\hbar} / \hbar = 0.1\%$

a) Calculate σ_E/E for arbitrary N .

b) If the precision of σ_E/E is to be improved which of the three quantities should be determined more precisely?

10) Taylor, Problem 3.28, page 85.

11) A Central Limit Theorem problem. When a certain chemical product is prepared the amount of a certain impurity is a random variable with a mean of 4 grams and a standard deviation of 2 grams. If 100 independent batches of the chemical are produced what is the (approximate) probability of the average amount of the impurity in the 100-batch sample being more than 4.5 grams?