

**K.K. Gan**  
**Physics 416**  
**Problem Set 2**

**Due Tuesday, April 26, 2010**

- 1) Assuming a Gaussian probability distribution answer the following questions

(Use Tables in *Taylor Appendix A and/or B*):

- a) What is the probability of a value lying more than  $1.5\sigma$  from the mean?
- b) What is the probability of a value lying  $\geq 1.5\sigma$  above the mean?
- c) What is the probability of a value lying  $\leq 1.5\sigma$  below the mean?
- d) What is the probability of a value,  $y$ , lying in the range  $\mu - \sigma \leq y \leq \mu + 2\sigma$ ?
- e) What is the probability of a value,  $y$ , lying in the range  $\mu + \sigma \leq y \leq \mu + 2\sigma$ ?

For this problem  $\mu$  is the mean of the Gaussian and  $\sigma$  is its standard deviation.

- 2) Taylor, Problem 5.12, page 156.

- 3) The sun emits an enormous number of neutrinos. Assume that  $10^6$  solar neutrinos uniformly pass through a square with an area of  $1 \text{ m}^2$  each  $\mu\text{sec}$ . Inside the square is a neutrino detector with an area of  $1 \text{ mm}^2$ . Assume Poisson statistics for this problem.

- a) What is the average number of neutrinos going through the detector each  $\mu\text{sec}$ ?
- b) What is the probability that no neutrinos go through the detector in a  $\mu\text{sec}$ ?
- c) What is the probability that  $\geq 2$  neutrinos go through the particle detector in a  $\mu\text{sec}$ ?
- d) How big should the detector be (in  $\text{mm}^2$ ) if we want  $\geq 2$  particles per  $\mu\text{sec}$  to pass through the detector with a probability of 95%?

- 4) Suppose a missile defense system destroys an incoming missile 95% of the time.

- a) If an evil country launches 20 missiles what is the probability that the missile defense system will destroy all of the incoming missiles?
- b) How many missiles have to be launched to have a 50% chance of at least one missile making it through the defense system?

Note: this problem can be done using either binomial or Poisson statistics.

- 5) According to quantum mechanics, the position ( $x$ ) of a particle in a one dimensional box with dimensions -  $L/2 \leq x \leq L/2$  ( $L$  constant) can be described by the following probability distribution function  $p(x)$ :

$$p(x) = A \cos^2[\pi x/L] \text{ for } -L/2 \leq x \leq L/2, \text{ and } 0 \text{ for all other } x.$$

- a) Find the normalization constant  $A$  in terms of  $L$ .
- b) Find the mean, mode, and median position of the particle in the box.
- c) Show that the variance ( $\sigma^2$ ) of  $x$  is given by:

$$\sigma^2 = \left(\frac{L}{\pi}\right)^2 \frac{\pi^2 - 6}{12}$$

- d) What is the probability of finding the particle in the region:  $L/4 \leq x \leq L/2$ ?

- 6) Suppose 100 six sided dice are tossed. Assume that the faces are labeled by one through six dots. Let  $Y_i$  be the number of dots on the  $i$ th ( $i=1$  to 100) die.

- a) What is the average number of dots expected for a single dice?
- b) What is the variance of the numbers of dots expected for a single dice?

c) Use the Central Limit Theorem to estimate the probability that the sum of the  $Y_i$ 's exceeds 400.

7) A Central Limit Theorem problem. When a certain chemical product is prepared the amount of a certain impurity is a random variable with a mean of 4 grams and a standard deviation of 2 grams. If 100 independent batches of the chemical are produced what is the (approximate) probability of the average amount of the impurity in the 100-batch sample being more than 4.5 grams?