

Lecture 8

Hypothesis Testing


Taylor Ch. 6 and 10.6

Introduction

- The goal of hypothesis testing is to set up a procedure(s) to allow us to decide if a mathematical model ("theory") is acceptable in light of our experimental observations.
- Examples:
 - ◆ Sometimes its easy to tell if the observations agree or disagree with the theory.
 - A certain theory says that Columbus will be destroyed by an earthquake in May 1992.
 - A certain theory says the sun goes around the earth.
 - A certain theory says that anti-particles (e.g. positron) should exist.
 - ◆ Often its not obvious if the outcome of an experiment agrees or disagrees with the expectations.
 - A theory predicts that a proton should weigh 1.67×10^{-27} kg, you measure 1.65×10^{-27} kg.
 - A theory predicts that a material should become a superconductor at 300K, you measure 280K.
 - ◆ Often we want to compare the outcomes of two experiments to check if they are consistent.
 - Experiment 1 measures proton mass to be 1.67×10^{-27} kg, experiment 2 measures 1.62×10^{-27} kg.

Types of Tests

- *Parametric Tests*: compare the values of parameters.
 - ◆ Example: Does the mass of the proton = mass of the electron?
- *Non-Parametric Tests*: compare the "shapes" of distributions.
 - ◆ Example: Consider the decay of a neutron. Suppose we have two theories that predict the energy spectrum of the electron emitted in the decay of the neutron (beta decay):



- Both theories might predict the same average energy for the electron.
 - ☞ A parametric test might not be sufficient to distinguish between the two theories.
- The shapes of their energy spectrums are quite different:
 - ★ Theory 1: the spectrum for a neutron decaying into two particles (e.g. $p + e$).
 - ★ Theory 2: the spectrum for a neutron decaying into three particles ($p + e + v$).
- ☞ We would like a test that uses our data to differentiate between these two theories.
- ◆ We can calculate the χ^2 of the distribution to see if our data was described by a certain theory:

$$\chi^2 = \sum_{i=1}^n \frac{(y_i - f(x_i, a, b))^2}{\sigma_i^2}$$

- $(y_i \pm \sigma_i, x_i)$ are the data points (n of them)
- $f(x_i, a, b, \dots)$ is a function that relates x and y
- ☞ accept or reject the theory based on the probability of observing a χ^2 larger than the above calculated χ^2 for the number of degrees of freedom.
- Example: We measure a bunch of data points $(y_i \pm \sigma_i, x_i)$ and we believe there is a linear relationship between x and y .

$$y = a + bx$$

- ★ If the y 's are described by a Gaussian PDF then minimizing the χ^2 function (or using LSQ or MLM method) gives an estimate for a and b .
- ★ As an illustration, assume that we have 6 data points and since we extracted a and b from the data, we have $6 - 2 = 4$ degrees of freedom (DOF). We further assume:

$$\chi^2 = \sum_{i=1}^6 \frac{(y_i - (a + bx_i))^2}{\sigma_i^2} = 15$$

- ☞ What can we say about our hypothesis that the data are described by a straight line?
- ☞ Look up the probability of getting $\chi^2 \geq 15$ by “chance”:

$$P(\chi \geq 15, 4) \approx 0.006$$
- ☞ only 6 of 1000 experiments would we expect to get this result ($\chi^2 \geq 15$) by “chance”.
- ☞ Since this is such a small probability we could reject the above hypothesis or we could accept the hypothesis and rationalize it by saying that we were unlucky.
- ☞ It is up to you to decide at what probability level you will accept/reject the hypothesis.

Confidence Levels (CL)

- An informal definition of a confidence level (CL):

$$CL = 100 \times [\text{probability of the event happening by chance}]$$
 - ◆ The 100 in the above formula allows CL's to be expressed as a percent (%).
- We can formally write for a continuous probability distribution P :

$$CL = 100 \times \text{prob}(x_1 \leq X \leq x_2) = 100 \times \int_{x_1}^{x_2} P(x) dx$$

For a CL, we know $P(x)$, x_1 , and x_2 .

- Example: Suppose we measure some quantity (X) and we know that X is described by a Gaussian distribution with mean $\mu = 0$ and standard deviation $\sigma = 1$.
 - ◆ What is the CL for measuring ≥ 2 (2σ from the mean)?

$$CL = 100 \times \text{prob}(X \geq 2) = 100 \times \frac{1}{\sigma\sqrt{2\pi}} \int_2^{\infty} e^{-\frac{(x-\mu)^2}{2\sigma^2}} dx = \frac{100}{\sqrt{2\pi}} \int_2^{\infty} e^{-\frac{x^2}{2}} dx = 2.5\%$$

- ◆ To do this problem we needed to know the underlying probability distribution P .
- ◆ If the probability distribution was not Gaussian (e.g. binomial) we could have a very different CL.
- ◆ If you don't know P you are out of luck!
- Interpretation of the CL can be easily abused.
 - ◆ Example: We have a scale of known accuracy (Gaussian with $\sigma = 10$ gm).
 - We weigh something to be 20 gm.
 - Is there really a 2.5% chance that our object weighs ≤ 0 gm??
 - ☞ probability distribution must be defined in the region where we are trying to extract information.

Confidence Intervals (CI)

- For a given confidence level, confidence intervals are the range $[x_1, x_2]$ that gives the confidence level.
 - ◆ Confidence interval's are not always uniquely defined.
 - ◆ We usually seek the minimum or symmetric interval.
- Example: Suppose we have a Gaussian distribution with $\mu = 3$ and $\sigma = 1$.
 - ◆ What is the 68% CI for an observation?
 - ◆ We need to find the limits of the integral $[x_1, x_2]$ that satisfy:

$$0.68 = \int_{x_1}^{x_2} P(x) dx$$

- ◆ For a Gaussian distribution the area enclosed by $\pm 1 \sigma$ is 0.68.

$$x_1 = \mu - 1 \sigma = 2$$

$$x_2 = \mu + 1 \sigma = 4$$

- ☞ confidence interval is $[2, 4]$.

For a CI, we know $P(x)$ and CL and wish to determine x_1 , and x_2 .

Upper/Lower Limits

- Example: Suppose an experiment observed no event.
 - ◆ What is the 90% CL upper limit on the expected number of events?

If $\lambda = 2.3$, then 10% of the time we expect to observe zero events even though there is nothing wrong with the experiment!

$$CL = 0.90 = \sum_{n=1}^{\infty} \frac{e^{-\lambda} \lambda^n}{n!}$$

$$1 - CL = 0.10 = 1 - \sum_{n=1}^{\infty} \frac{e^{-\lambda} \lambda^n}{n!} = \sum_{n=0}^{\infty} \frac{e^{-\lambda} \lambda^n}{n!} = e^{-\lambda}$$

$$\lambda = 2.3$$

- ◆ If the expected number of events is greater than 2.3 events,
 - ☞ the probability of observing one or more events is greater than 90%.
- Example: Suppose an experiment observed one event.
 - ◆ What is the 95% CL upper limit on the expected number of events?

$$CL = 0.95 = \sum_{n=2}^{\infty} \frac{e^{-\lambda} \lambda^n}{n!}$$

$$1 - CL = 0.05 = 1 - \sum_{n=2}^{\infty} \frac{e^{-\lambda} \lambda^n}{n!} = \sum_{n=0}^1 \frac{e^{-\lambda} \lambda^n}{n!} = e^{-\lambda} + \lambda e^{-\lambda}$$

$$\lambda = 4.74$$

Procedure for Hypothesis Testing

- a) Measure something.
 - b) Get a hypothesis (sometimes a theory) to test against your measurement.
 - c) Calculate the CL that the measurement is from the theory.
 - d) Accept or reject the hypothesis (or measurement) depending on some minimum acceptable CL.
- Problem: How do we decide what is acceptable CL?
 - ◆ Example: What is an acceptable definition that the space shuttle is safe?
 - ★ One explosion per 10 launches or per 1000 launches or...?

Hypothesis Testing for Gaussian Variables

- If we want to test whether the mean of some quantity we have measured (x = average from n measurements) is consistent with a known mean (μ_0) we have the following two tests:

Test	Condition	Test Statistic	Test Distribution
$\mu = \mu_0$	σ^2 known	$\frac{x - \mu_0}{\sigma/\sqrt{n}}$	Gaussian $\mu = 0, \sigma = 1$
$\mu = \mu_0$	σ^2 unknown	$\frac{x - \mu_0}{s/\sqrt{n}}$	$t(n - 1)$

- ◆ s : standard deviation extracted from the n measurements.
- ◆ $t(n - 1)$: Student's "t-distribution" with $n - 1$ degrees of freedom.
 - Student is the pseudonym of statistician W.S. Gosset who was employed by a famous English brewery.
- Example: Do free quarks exist? Quarks are nature's fundamental building blocks and are thought to have electric charge (q) of either $(1/3)e$ or $(2/3)e$ (e = charge of electron). Suppose we do an experiment to look for $q = 1/3$ quarks.
 - ◆ Measure: $q = 0.90 \pm 0.2 = \mu \pm \sigma$
 - ◆ Quark theory: $q = 0.33 = \mu_0$
 - ◆ Test the hypothesis $\mu = \mu_0$ when σ is known:
 - ☞ Use the first line in the table:

$$z = \frac{x - \mu_0}{\frac{\sigma}{\sqrt{n}}} = \frac{0.9 - 0.33}{\frac{0.2}{\sqrt{1}}} = 2.85$$

- Assuming a Gaussian distribution, the probability for getting a $z \geq 2.85$,

$$\text{prob}(z \geq 2.85) = \int_{2.85}^{\infty} P(\mu, \sigma, x) dx = \int_{2.85}^{\infty} P(0, 1, x) dx = \frac{1}{\sqrt{2\pi}} \int_{2.85}^{\infty} e^{-\frac{x^2}{2}} dx = 0.002$$

- If we repeated our experiment 1000 times,
 - ☞ two experiments would measure a value $q \geq 0.9$ if the true mean was $q = 1/3$.
 - ☞ This is not strong evidence for $q = 1/3$ quarks!
- ◆ If instead of $q = 1/3$ quarks we tested for $q = 2/3$ what would we get for the CL?
 - $\mu = 0.9$ and $\sigma = 0.2$ as before but $\mu_0 = 2/3$.
 - ☞ $z = 1.17$
 - ☞ $\text{prob}(z \geq 1.17) = 0.13$ and $\text{CL} = 13\%$.
 - ☞ quarks are starting to get believable!
- Consider another variation of $q = 1/3$ problem. Suppose we have 3 measurements of the charge q : $q_1 = 1.1$, $q_2 = 0.7$, and $q_3 = 0.9$
 - ◆ We don't know the variance beforehand so we must determine the variance from our data.
 - ☞ use the second test in the table:

$$\mu = \frac{1}{3}(q_1 + q_2 + q_3) = 0.9$$

$$s^2 = \frac{\sum_{i=1}^n (q_i - \mu)^2}{n - 1} = \frac{0.2^2 + (-0.2)^2 + 0}{2} = 0.04$$

$$z = \frac{x - \mu_0}{s/\sqrt{n}} = \frac{0.9 - 0.33}{0.2/\sqrt{3}} = 4.94$$
 - Table 7.2 of Barlow: $\text{prob}(z \geq 4.94) \approx 0.02$ for $n - 1 = 2$.
 - ☞ 10X greater than the first part of this example where we knew the variance ahead of time.
- Consider the situation where we have several independent experiments that measure the same quantity:
 - ◆ We do not know the true value of the quantity being measured.
 - ◆ We wish to know if the experiments are consistent with each other.

Test	Conditions	Test Statistic	Test Distribution
$\mu_1 = \mu_2$	σ_1^2 and σ_2^2 known	$\frac{x_1 - x_2}{\sqrt{\sigma_1^2/n + \sigma_2^2/m}}$	Gaussian $\mu = 0, \sigma = 1$
$\mu_1 = \mu_2$	$\sigma_1^2 = \sigma_2^2 = \sigma^2$ unknown	$\frac{x_1 - x_2}{Q\sqrt{1/n + 1/m}}$	$t(n + m - 2)$
$\mu_1 = \mu_2$	$\sigma_1^2 \neq \sigma_2^2$ unknown	$\frac{x_1 - x_2}{\sqrt{s_1^2/n + s_2^2/m}}$	approx. Gaussian $\mu = 0, \sigma = 1$

$$Q^2 \equiv \frac{(n-1)s_1^2 + (m-1)s_2^2}{n+m-2}$$

- Example: We compare results of two independent experiments to see if they agree with each other.

Exp. 1 1.00 ± 0.01

Exp. 2 1.04 ± 0.02

- Use the first line of the table and set $n = m = 1$.

$$z = \frac{x_1 - x_2}{\sqrt{\sigma_1^2/n + \sigma_2^2/m}} = \frac{1.04 - 1.00}{\sqrt{0.01^2 + 0.02^2}} = 1.79$$

- z is distributed according to a Gaussian with $\mu = 0, \sigma = 1$.
- Probability for the two experiments to disagree by ≥ 0.04 :

$$\text{prob}(|z| \geq 1.79) = 1 - \int_{-1.79}^{1.79} P(\mu, \sigma, x) dx = 1 - \int_{-1.79}^{1.79} P(0, 1, x) dx = 1 - \frac{1}{\sqrt{2\pi}} \int_{-1.79}^{1.79} e^{-\frac{x^2}{2}} dx = 0.07$$

- ★ We don't care which experiment has the larger result so we use $\pm z$.
- 7% of the time we should expect the experiments to disagree at this level.
- Is this acceptable agreement?