

Supplementary Information for: “Structure of the aqueous electron”

John M. Herbert*

Department of Chemistry & Biochemistry
The Ohio State University, Columbus, OH USA

August 30, 2019

*herbert@chemistry.ohio-state.edu




Figure S1: Radial distribution functions (RDFs), $g(r)$, for the Larsen-Glover-Schwartz (LGS),¹ Turi-Borgis (TB),^{2,3} and Jacobson-Herbert⁴ (PEWP-2) pseudopotential models of $e^-(aq)$. Panel (a) shows the $e^- \cdots$ hydrogen RDF and (b) shows the $e^- \cdots$ oxygen RDF, where the electron coordinate (e^-) is the centroid of the one-electron wave function. These plots were constructed using simulation data originally reported in Ref. 5.




Figure S2: Overlay of the RDFs from Fig. 7, representing $g(r)$ data from from two different QM/MM simulations of $e^-(aq)$, one at the BLYP(SIC)+D3 level (from Ref. 6) and the other at the HF+D3 level (from Ref. 7). The RDFs for the $e^- \cdots$ H coordinate are shown in green [BLYP(SIC)+D3] and in orange (HF+D3), while those for the $e^- \cdots$ O coordinate are shown in red [BLYP(SIC)+D3] and in blue (HF+D3). Dashed curves are the integrated coordination numbers $n(r)$ from the BLYP(SIC)+D3 simulation and should be read from the axis on the right. The green and red arrows that point to the axis on the right suggest coordination numbers obtained by integrating Eq. (4) that defines $n(r)$, up to the first local minimum in either RDF.




Figure S3: Resonance Raman (RR) spectra of e^- (aq) computed within the instantaneous normal modes (INM) approach, using the excited-state gradient approximation of Eq. (6).⁸ (a) INM-RR spectrum computed at non-cavity liquid geometries obtained from a simulation using the LGS model. (b) INM-RR spectrum computed from a QM/MM trajectory at the HF+D3/3-21++G* level, which supports a stable cavity. These spectra represent ensemble averages of harmonic stick spectra computed for QM/MM snapshots and then broadening with 20 cm^{-1} Lorentzian functions, with RR intensity enhancements computed according to Eq. (6). The INM-RR spectra in Figs. 8 and 9 were computed in the same way but use Lorentzian broadening of 45 cm^{-1} .

Adapted from Ref. 8; copyright 2019 American Chemical Society.




Figure S4: Temperature dependence of the aqueous electron's absorption maximum, $E_{\max}(T)$. (a) Comparison of predictions from the TB, SR, LGS, and PEWP-2 pseudopotential models. All simulation data are from Ref. 9 except for the PEWP-2 data, which are new.¹⁰ All simulations were carried out at a fixed water density of 0.997 g/cm^3 . The SR and TB values of E_{\max} are shifted by -0.50 eV and -0.25 eV , respectively, as indicated. Experimental data are from Ref. 11. (b) Predictions from the TB model using variable densities, where the density used in the simulation corresponds to the experimental density of neat liquid water at the indicated temperature. Results plotted are from simulations reported in Ref. 12 and experiments reported in Ref. 13. Note that the energy (vertical) scale is the same in both panels but that the data in (b) span a much broader range of temperatures, including supercritical data.




Figure S5: Absorption spectra computed using the Jacobson-Herbert (PEWP-2) pseudopotential model at several different temperatures, using bin widths of (a) 0.10 eV and (b) 0.15 eV . Excitation energy data up to 2.5 eV were fit to a Gaussian envelope, with E_{\max} as one fitting parameter. This value of E_{\max} is then used to plot the T -dependent absorption maximum for the PEWP-2 model in Fig. S4.

References

- [1] R. E. Larsen, W. J. Glover, and B. J. Schwartz, Does the hydrated electron occupy a cavity?, *Science*, 2010, **329**, 65–69.
- [2] L. Turi, M.-P. Gaigeot, N. Levy, and D. Borgis, Analytical investigations of an electron–water molecule pseudopotential. I. Exact calculations on a model system, *J. Chem. Phys.*, 2001, **114**, 7805–7815.
- [3] L. Turi and D. Borgis, Analytical investigations of an electron–water molecule pseudopotential. II. Development of a new pair potential and molecular dynamics simulations, *J. Chem. Phys.*, 2002, **117**, 6186–6195.
- [4] L. D. Jacobson and J. M. Herbert, A one-electron model for the aqueous electron that includes many-body electron-water polarization: Bulk equilibrium structure, vertical electron binding energy, and optical absorption spectrum, *J. Chem. Phys.*, 2010, **133**, 154506:1–19.
- [5] J. M. Herbert and L. D. Jacobson, Structure of the aqueous electron: Assessment of one-electron pseudopotential models in comparison to experimental data and time-dependent density functional theory, *J. Phys. Chem. A*, 2011, **115**, 14470–14483.
- [6] F. Uhlig, O. Marsalek, and P. Jungwirth, Unraveling the complex nature of the hydrated electron, *J. Phys. Chem. Lett.*, 2012, **3**, 3071–3075, Erratum: *ibid.* **4**, 603 (2013).
- [7] Z. C. Holden, B. Rana, and J. M. Herbert, Analytic energy gradients for the QM/MM-Ewald method using atomic charges derived from the electrostatic potential: Theory, implementation, and application to *ab initio* molecular dynamics of the aqueous electron, *J. Chem. Phys.*, 2019, **150**, 144115:1–20.
- [8] S. Dasgupta, B. Rana, and J. M. Herbert, *Ab initio* investigation of the resonance Raman spectrum of the hydrated electron, 2019 (in press; DOI: 10.1021/acs.jpcc.9b04895).
- [9] J. R. Casey, R. E. Larsen, and B. J. Schwartz, Resonance Raman and temperature-dependent electronic absorption spectra of cavity and noncavity models of the hydrated electron, *Proc. Natl. Acad. Sci. USA*, 2013, **110**, 2712–2717.
- [10] The PEWP-2 simulations were performed with 200 molecules in a periodic simulation cell of dimensions $L \times L \times L$, where $L = 18.1671 \text{ \AA}$ corresponding to a water density of 0.997 g/cm^3 . The same water density is used at each temperature, with temperature conservation (NVT ensemble) enforced by means of a Nosé-Hoover thermostat. Other simulation details follow the calculations reported in Ref. 4.
- [11] Y. Du, E. Price, and D. M. Bartels, Solvated electron spectrum in supercooled water and ice, *Chem. Phys. Lett.*, 2007, **438**, 234–237.
- [12] C. Nicolas, A. Boutin, B. Levy, and D. Borgis, Molecular simulation of a hydrated electron at different thermodynamic state points, *J. Chem. Phys.*, 2003, **118**, 9689–9696.
- [13] G. Wu, Y. Katsumura, Y. Muroya, X. Li, and Y. Terada, Hydrated electron in subcritical and supercritical water: A pulse radiolysis study, *Chem. Phys. Lett.*, 2000, **325**, 531–536.