


Homework Set No. 4, Physics 880.08

Deadline – noon on Monday, March 12, 2012

1. (15 pts) Show that the light-by-light scattering QED diagram pictured below does not contain a UV divergence (even though naively counting powers of momenta makes it seem logarithmically divergent). The ellipsis in the figure represent all possible permutations of the vertices along the loop. (Hint: you may find formulas (A.41) and (A.42) in Appendix A.4 of Peskin and Schroeder useful.)



2. Consider φ^4 scalar theory with the Lagrangian

$$\mathcal{L} = \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi - \frac{m^2}{2} \varphi^2 - \frac{\lambda}{4!} \varphi^4.$$

Below use Pauli-Villars regularization with two new particles, such that the propagators of the internal lines are replaced by

$$\frac{i}{k^2 - m^2 + i\epsilon} \rightarrow \frac{i}{k^2 - m^2 + i\epsilon} + \sum_{j=1}^2 \frac{i C_j}{k^2 - M_j^2 + i\epsilon}$$

with $C_1 = 1$, $C_2 = -2$, and $M_1^2 = m^2 + 2M^2$, $M_2^2 = m^2 + M^2$. M is some UV cutoff.

a. (30 pts) Calculate regularized one-loop 1PI corrections to the propagator and to the four-point Green function, keeping only divergent and finite terms at large M (i.e., dropping all terms which vanish in the $M \rightarrow \infty$ limit).

b. (15 pts) Set up the renormalization program for the ϕ^4 theory at one loop: that is, find all counterterms and coefficients in front of them using “on-shell” renormalization

conditions:

$$\begin{aligned}\Sigma(p^2 = m^2) &= 0, \\ \left. \frac{\partial \Sigma(p^2)}{\partial p^2} \right|_{p^2=m^2} &= 0, \\ \Gamma^4(s = 4m^2, t = 0, u = 0) &= -i\lambda.\end{aligned}$$

Here $-i\Sigma(p^2)$ is the sum of all 1PI corrections to the propagator, and $\Gamma^4(s, t, u)$ is the truncated 1PI 4-point Green function. Also m is the physical particle mass, and λ is the physical coupling.

In particular show that the renormalized 4-point function $\Gamma_R^4(s, t, u)$ is given by

$$\begin{aligned}\Gamma_R^4(s, t, u) = -i \left\{ \lambda + \frac{\lambda^2}{32\pi^2} \int_0^1 dx \left[\ln \left(\frac{m^2 - sx(1-x)}{m^2 - 4m^2x(1-x)} \right) + \ln \left(\frac{m^2 - tx(1-x)}{m^2} \right) \right. \right. \\ \left. \left. + \ln \left(\frac{m^2 - ux(1-x)}{m^2} \right) \right] + o(\lambda^3) \right\}.\end{aligned}$$

For a 4-point function with two external lines having incoming 4-momenta p_1^μ and p_2^μ and two other external lines having outgoing 4-momenta p_3^μ and p_4^μ we define as usual $s = (p_1 + p_2)^2$, $t = (p_1 - p_3)^2$ and $u = (p_1 - p_4)^2$. (Hint: you may find the discussion in Chapter 10.2 of Peskin and Schroeder useful, though they use dimensional regularization.)