

EXTENSION OF THE MEASUREMENT, ASSIGNMENT, AND FIT IN THE GROUND STATE OF THE TWO-TOP MOLECULE METHYL ACETATE

H. V. L. NGUYEN, I. KLEINER, *Laboratoire Interuniversitaire des Systèmes Atmosphériques (LISA), UMR 7583 (CNRS/Univ. Paris Est et Paris Diderot), Université de Paris Est, 61 avenue du Général de Gaulle, F-94010 Créteil cedex, France*; S. SHIPMAN, *Division of Natural Sciences, New College of Florida, 5800 Bay Shore Road, Sarasota, FL 34243-2109, USA*; and K. KOBAYASHI, *Department of Physics, Faculty of Science, Toyama University, 3190 Gofuku Toyama, Toyama, 930-8555, Japan*.

New spectroscopic data were recorded for the two-top molecule methyl acetate with a chirped-pulse Fourier transform microwave spectrometer at room temperature and a jet cooled sub-millimeterwave spectrometer. More than 650 new lines with J up to 35 and K up to 15 were assigned. In total, approximately 1500 lines were fitted with 34 parameters using the program BELGI-C_s-2tops to a standard deviation close to the measurement error. More precise determinations of the top-top interaction and the J , K dependent parameters as well as the influence of the total rotation on the top-top interaction were carried out. The extension of the ground state spectrum of methyl acetate is a necessary step before searching for the first torsional excited transition in both torsional modes in the room temperature spectrum.