

DISPERSED FLUORESCENCE SPECTROSCOPY OF THE $\tilde{B}^2E' - \tilde{X}^2A'_2$ TRANSITION OF JET COOLED $^{14}\text{NO}_3$ and $^{15}\text{NO}_3$

MASARU FUKUSHIMA and TAKASHI ISHIWATA, *Faculty of Information Sciences, Hiroshima City University, Asa-Minami, Hiroshima 731-3194, Japan.*

We have generated NO_3 in supersonic free jet expansions and observed laser induced fluorescence (LIF) of the $\tilde{B}^2E' - \tilde{X}^2A'_2$ transition. We have measured LIF excitation spectra and dispersed fluorescence (DF) spectra from the single vibronic levels (SVL's) of the \tilde{B}^2E' state of $^{14}\text{NO}_3$ and $^{15}\text{NO}_3$. The vibrational structure of the $\tilde{X}^2A'_2$ state has been analyzed by comparing the vibrational structures of the DF spectra of the two isotopomers. The $1,053\text{ cm}^{-1}$ band of $^{14}\text{NO}_3$ is observed as two bands at $1,039$ and $1,053\text{ cm}^{-1}$ with an intensity ratio of $4 : 5$, respectively, for $^{15}\text{NO}_3$, which are observed in the DF spectra with our standard resolution ($\sim 7\text{ cm}^{-1}$ in FWHM). Higher resolution measurements ($\sim 2\text{ cm}^{-1}$ in FWHM) of the DF spectra show that the $1,053\text{ cm}^{-1}$ band of $^{14}\text{NO}_3$ is also observed as two bands at $1,051$ and $1,056\text{ cm}^{-1}$ with an intensity ratio of $5 : 3$, respectively. The $1,051\text{ cm}^{-1}$ band is attributed to be the ν_1 (a_1') fundamental, because of its little isotope shift. There are two possibilities for another band, the band at $1,056$ and $1,038\text{ cm}^{-1}$ for $^{14}\text{NO}_3$ and $^{15}\text{NO}_3$, respectively; (1) the ν_3 (e') fundamental band^a, and (2) the $\nu_2 + \nu_4$ (a_2'' and e' , respectively) combination band. If this is the case (1), the ν_3 band should be observed in IR spectrum, but it has yet to be observed. If (2), the intensity must be stolen from the $\tilde{B}^2E' - \tilde{A}^2E''$ transition through the ν_2 mode, the considerable transition moment of which has been predicted^b. A simple consideration for the vibronic coupling^c between the \tilde{A}^2E'' and $\tilde{X}^2A'_2$ states through the ν_2 mode can understand about 20 % of the combination band intensity to that of the ν_1 fundamental. The higher resolution measurements of the DF spectra also show that the $1,499\text{ cm}^{-1}$ band of $^{14}\text{NO}_3$ is much stronger than the $1,492\text{ cm}^{-1}$ band in the electronic spectrum, while the latter is the strongest band in the IR absorption spectrum^d.

^aJ. F. Stanton, J. Chem. Phys. 126, 134309 (2007).

^bJ. F. Stanton and M. Okumura, Phys. Chem. Chem. Phys. 11, 4742 (2009).

^cE. Hirota, K. Kawaguchi, T. Ishiwata, and I. Tanaka, J. Chem. Phys. 95, 771 (1991).

^dT. Ishiwata, I. Tanaka, K. Kawaguchi, and E. Hirota, J. Chem. Phys. 82, 2196 (1985).