

## ROTATIONALLY RESOLVED HIGH-RESOLUTION LASER SPECTROSCOPY OF THE $S_1 \leftarrow S_0$ TRANSITION OF NAPHTHALENE AND Cl-NAPHTHALENE

SHUNJI KASAHARA, RYO YAMAMOTO, KOHEI TADA, *Molecular Photoscience Research Center, Kobe University, Kobe 657-8501, Japan.*

Rotationally resolved high-resolution fluorescence excitation spectra and the Zeeman effects of 0-0 band of  $S_1 \leftarrow S_0$  electronic transition have been observed for naphthalene, 1-Cl naphthalene (1-CIN), and 2-Cl naphthalene (2-CIN). Sub-Doppler excitation spectra were measured by crossing a single-mode UV laser beam perpendicular to a collimated molecular beam. The typical linewidth was 25 MHz and the absolute wavenumber was calibrated with accuracy  $0.0002 \text{ cm}^{-1}$  by measurement of the Doppler-free saturation spectrum of iodine molecule and fringe pattern of the stabilized etalon. For naphthalene<sup>a</sup> and 2-CIN, the rotationally resolved spectra were obtained, and these molecular constants were determined in high accuracy. The obtained molecular constants of 2-CIN are in good agreement with the ones reported by Plusquellic *et. al.*<sup>b</sup> For 1-CIN, the rotational lines were not completely resolved because the fluorescence lifetime is shorter than the one of 2-CIN. Additionally, we have observed the change of the spectra with magnetic field. The Zeeman broadening was mainly observed for the levels of low  $K_a$  and increasing in proportion to  $J$  for given  $K$  for both of naphthalene and 2-CIN. The order of magnitude and the  $J, K$ -dependence of the observed Zeeman broadening were similar to the other vibronic bands of naphthalene.<sup>c</sup>

---

<sup>a</sup>D. L. Joo, R. Takahashi, J. O'Reilly, H. Katô, and M. Baba, *J. Mol. Spectrosc.*, **215**, 155 (2002).

<sup>b</sup>D. F. Plusquellic, S. R. Davis, and F. Jahanmir, *J. Chem. Phys.*, **115**, 225 (2001).

<sup>c</sup>H. Katô, S. Kasahara, and M. Baba, *Bull. Chem. Soc. Jpn.*, **80**, 456 (2007).