

TERAHERTZ SPECTROSCOPY OF CaH ($X^2\Sigma^+$), MgH ($X^2\Sigma^+$), AND ZnH ($X^2\Sigma^+$) : EXTREME HYDRIDE SYNTHESIS

MATTHEW P. BUCCHINO, *Department of Chemistry and Biochemistry, University of Arizona, Tucson, Arizona 85721*; and LUCY M. ZIURYS, *Department of Chemistry and Biochemistry, Department of Astronomy, and Steward Observatory, University of Arizona, Tucson, Arizona 85721*.

Sub-millimeter spectra of CaH ($X^2\Sigma^+$), MgH ($X^2\Sigma^+$), and ZnH ($X^2\Sigma^+$) have been measured in the sub-mm/THz regime (500–805 GHz) using direct-absorption methods. All species were produced by the reaction of metal vapor with H₂ under DC discharge in a Broida-type oven. Because of efficient molecule production, spectra of ²⁶MgH, ⁶⁶ZnH, ⁶⁸ZnH and ⁷⁰ZnH could be recorded in their natural abundance as well, with high signal-to-noise. Additionally, the N = 0 → 1 and N = 1 → 2 transitions of ²⁵MgH and ⁶⁷ZnH were measured. Fermi contact, dipolar, and quadrupole constants for the ²⁵Mg and ⁶⁷Zn nuclei have been accurately determined for the first time in the gas phase, and are in good agreement with previous ESR studies. The hyperfine constants suggest a substantial amount of covalent character is present in metal hydrides.