

HIGH-RESOLUTION PHOTOELECTRON SPECTROSCOPY OF 2-BUTYNE

UGO JACOVELLA, BÉRENGER GANS and FRÉDÉRIC MERKT, *ETH Zürich, Laboratorium für Physikalische Chemie, Wolfgang-Pauli-Strasse 10, 8093 Zürich, Switzerland.*

Using a coherent narrow-band vacuum-ultraviolet (VUV) laser source (bandwidth of 0.008 cm^{-1})^a coupled to a photoionization and pulse-field-ionization zero-kinetic-energy photoelectron (PFI-ZEKE) spectrometer, the threshold photoionization of polyatomic molecules can be studied at high resolution.

We present a new measurement of the PFI-ZEKE photoelectron spectrum of the origin band of the $X^+ \ ^2E_{2(d)} \leftarrow X \ ^1A_{1(s)}$ ionizing transition of 2-butyne at a resolution of 0.15 cm^{-1} . Despite this high resolution, the spectral congestion originating from the combined effects of the internal rotation, the spin-orbit coupling and the Jahn-Teller effect prevented the full resolution of the rotational structure of the photoelectron spectrum. Combined with the known structure of the $X \ ^1A_{1(s)}$ ground state of 2-butyne, including the free internal rotation^{bcd}, the spectrum was used to derive information on the $X^+ \ ^2E_{2(d)}$ ground state of the 2-butyne radical cation. The rotational branch structure of the spectrum points at a complex energy-level structure of the cation and at the importance of a shape resonance enhancing g photoelectron partial waves^e.

^aU. Hollenstein, H. Palm, and F. Merkt, *Rev. Sci. Instrum.* **71**, 4023 (2000).

^bH. C. Longuet-Higgins *Mol. Phys.* **6**, 445 (1963).

^cJ. T. Hougen *J. Chem. Phys.* **37**, 1433 (1962).

^dP. R. Bunker *Mol. Phys.* **8**, 81 (1964).

^eH. Xu, U. Jacovella, B. Ruscic, S. T. Pratt and R. R. Lucchese *J. Chem. Phys.* **136**, 154303 (2012).