

## DETECTION OF NO AND S-NITROSO COMPOUNDS USING MID-IR CAVITY RING-DOWN SPECTROSCOPY

VITALI STSIAPURA, VINCENT K. SHUALI, ANGELA ZIEGLER, KEVIN K. LEHMANN, *Chemistry Department, University of Virginia, Charlottesville, VA 22904*; and BENJAMIN M. GASTON, *Case Western Reserve University, Cleveland, Ohio 44106*.

Metabolic reactions of S-nitrosothiols have received much attention in biochemistry and medicine since S-nitroso compounds can act not only as donors of nitric oxide (NO) in an organism but can themselves be involved in signal transduction. In the past few decades, the primary means of detecting S-nitroso compounds in biological media has been through chemiluminescence detection of NO, a technique that limits the sensitivity to ppb levels and is not able to trace isotopologues of NO.

Here we present a cw-CRD instrument for detection of NO isotopologues released from S-nitroso compounds with a sensitivity of 27 pptv of NO in 100 torr He (after averaging of  $\sim 400$  ringdowns). A mid-IR ec-QCL laser (Daylight Solutions) is used to excite the ringdown cavity of finesse  $\sim 10000$  and probe the NO fundamental ro-vibrational band in the 5.2 - 5.3  $\mu\text{m}$  range.