

PRECISION LASER SPECTROSCOPY OF H_3^+

HSUAN-CHEN CHEN, *Institute of Photonics Technologies, National Tsing Hua University, Hsinchu 30013, Taiwan*; JIN-LONG PENG, *Center for Measurement Standards, Industrial Technology Research Institute, Hsinchu 30011, Taiwan*; T. AMANO, *Department of Chemistry and Department of Physics and Astronomy, University of Waterloo, Waterloo, ON N2L 3G1, Canada*; JOW-TSONG SHY, *Institute of Photonics Technologies, National Tsing Hua University and Department of Physics, National Tsing Hua University, Hsinchu 30013, Taiwan*.

The high-resolution sub-Doppler Lamb dips of the ν_2 fundamental band transitions of H_3^+ have been observed using an extended negative glow discharge tube as an ion source and a periodically poled lithium niobate optical parametric oscillator as a radiation source.^a The absolute frequencies of five transitions were measured to an accuracy of 250 kHz using a fiber optical frequency comb. In addition, we measured the homogeneous linewidths of these lines. Physical significance of these results will be discussed in terms of collisional processes.

^aH.-C. Chen, C.-Y. Hsiao, J.-L. Peng, T. Amano, and J.-T. Shy, *Phys. Rev. Lett.*, **109**, 263002 (2012)