

ROTATIONAL SPECTRA OF ADDUCTS OF FORMALDEHYDE WITH FREONS

GOU QIAN, GANG FENG, LUCA EVANGELISTI, and W. CAMINATI, *Dipartimento di Chimica "G. Ciamician" dell'Università, Via Selmi 2, I-40126 Bologna, Italy*; MONTSERRAT VALLEJO LÓPEZ, ALBERTO LESARRI, *Departamento de Química Física y Química Inorgánica, Facultad de Ciencias, Universidad de Valladolid, E-47011 Valladolid, Spain*; EMILIO COCINERO, *Departamento de Química Física, Facultad de Ciencia y Tecnología, Universidad del País Vasco, E-48080 Bilbao, Spain*.

The rotational spectra of three 1:1 complexes of formaldehyde (H_2CO) with freons, i.e. difluoromethane (CH_2F_2), fluorochloromethane (CH_2FCl) and trifluorochloromethane (CF_3Cl), have been observed and assigned using pulsed jet Fourier transform microwave technique. Several isotopologues (including some ^{13}C species) have been measured in natural abundance. The tunnelling splittings have been measured in the first two adducts with relative intensity 1:3, due to the internal rotation of the formaldehyde moiety along its symmetry axis. The barriers to this motion have been estimated by using a flexible model. For the latter two complexes, each of transition displays the hyperfine structures due to the quadrupolar effects of ^{35}Cl (^{37}Cl) nucleus. The dissociation energy has been estimated within the pseudo-diatomic approximation for all three complexes.