

FELION: A CRYOGENIC ION TRAP APPARATUS FOR SPECTROSCOPIC STUDIES WITH FELIX

S. BRÜNKEN, L. KLUGE, S. FANGHÄNEL, A. POTAPOV, O. ASVANY, and S. SCHLEMMER, *I. Physikalisches Institut, Universität zu Köln, 50937 Köln, Germany*; J. OOMENS, B. REDLICH, *Radboud University Nijmegen, Institute for Molecules and Materials, FELIX Facility, 6525 ED Nijmegen, Netherlands*; A. STOFFELS, *I. Physikalisches Institut, Universität zu Köln, 50937 Köln, Germany, and Radboud University Nijmegen, Institute for Molecules and Materials, FELIX Facility, 6525 ED Nijmegen, Netherlands.*

The combination of ion trapping techniques with sensitive action spectroscopy schemes has been developed in recent years as a powerful tool to obtain spectra of gas-phase molecular ions from the UV to the (F)IR spectral regions.^a Here we report on the status of a specifically designed, dedicated cryogenic 22-pole ion trap apparatus (FELion), developed and built in Cologne, Germany, and intended to be installed permanently at the "Free-Electron Lasers for Infrared eXperiments" (FELIX) facility in Nijmegen, the Netherlands. This instrument will allow to record gas-phase IR and FIR spectra of mass-selected, internally cold molecular ions at temperatures in the range 4 – 300 K. By the use of diverse ionization methods, e.g. electron impact and electrospray ionization, a multitude of molecular ions can be generated and stored in the trap, including astrophysically relevant species ranging in size from the three-atomic H_3^+ up to large polycyclic aromatic hydrocarbon (PAH) ions, but also biomolecular ions like amino acids, peptides, or nucleobases. In combination with the powerful (F)IR radiation of the free electron lasers FELIX-1 and -2 ($60 - 2500 \text{ cm}^{-1}$) and FLARE^b ($6 - 100 \text{ cm}^{-1}$) at the FELIX facility, a variety of action spectroscopy schemes can be employed to study the ro-vibrational spectra of the stored ions, such as IR multiphoton dissociation, (F)IR/UV double resonance spectroscopy, or the method of laser induced reactions (LIR). In this talk we will give a detailed account of the experimental setup and present the first results obtained with the new apparatus.

^ae.g., S. Schlemmer, E. Lescop, J. von Richthofen, D. Gerlich, and M. A. Smith, *J. Chem. Phys.* 117, 2068 (2002); J. Oomens, B. G. Sartakov, G. Meijer, and G. van Helden, *Int. J. Mass Spectrom.* 254, 1 (2006); T. R. Rizzo, J. A. Stearns, and O. V. Boyarkin, *International Reviews in Physical Chemistry* 28, 481 (2009)

^bFLARE: Free-electron Laser for Advanced spectroscopy and high-Resolution Experiments