

FREQUENCY-COMB REFERENCED, SUB-DOPPLER SPECTROSCOPY OF HOT BANDS OF ACETYLENE IN THE REGION OF THE $\nu_1 + \nu_3$ COMBINATION BAND

CHRISTOPHER P. MCRAVEN, *Chemistry Department, Brookhaven National Laboratory, Upton, New York 11973*; MATTHEW J. CICH, SALVATORE M. CAIOLA, *Department of Chemistry, Stony Brook University, Stony Brook, New York 11794*; TREVOR J. SEARS^a, *Chemistry Department, Brookhaven National Laboratory, Upton, New York 11973*; GREGORY E. HALL, *Chemistry Department, Brookhaven National Laboratory, Upton, New York 11973*.

To take full advantage of recent improvements in the accuracy obtained in collision-induced absorption line shape measurements of the acetylene $\nu_1 + \nu_3$ combination band, independent and accurate knowledge of the positions of weaker, overlapping hot band transitions is required. We have used cavity-enhanced saturation-absorption spectroscopy together with a frequency-comb referenced, extended cavity diode laser to determine Doppler-free transition frequencies for the ν_4 and ν_5 hot band transitions in this region. A laser intensity dependent asymmetry in the sub-Doppler saturation line shape is observed, and will be discussed in terms of the intra-cavity beam geometry.

Acknowledgments: Work at Brookhaven National Laboratory was carried out under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy and supported by its Office of Basic Energy Sciences, Division of Chemical Sciences, Geosciences and Biosciences.

^aalso: *Department of Chemistry, Stony Brook University, Stony Brook, New York 11794*