

## PURE ROTATIONAL SPECTRA OF THE REACTION PRODUCTS OF LASER ABLATED THORIUM METAL AND OXYGEN MOLECULES ENTRAINED WITHIN SUPERSONIC EXPANSIONS OF NOBLE GASES

B. E. LONG, *Department of Chemistry, Wesleyan University, 52 Lawn Avenue, Middletown, Connecticut, 06459*; S. A. COOKE, *School of Natural and Social Sciences, Purchase College SUNY, 735 Anderson Hill Road, Purchase, NY 10577, USA*.

Thorium metal has been laser ablated in the presence of pure oxygen entrained in high pressure argon. The products of the ablation event have been supersonically expanded into the Fabry-Pérot cavity of a time-domain microwave spectrometer. New measurements have been performed on thorium monoxide, where the  $J = 1 \leftarrow 0$  transition has been recorded in the  $v = 8, 9,$  and  $10$  vibrational levels. Further to this, a thorium-dependent spectral transition has been observed at  $19251.8740(10)$  MHz. The carrier of this signal is to be determined, however, it requires an oxygen concentration of  $0.1\%$  which is significantly higher than the optimal oxygen concentration required for observation of the thorium monoxide transitions. It is postulated that the presently unidentified thorium-containing compound is  $\text{ThO}_2$  which has  $C_{2v}$  symmetry. Progress on the measurement of this spectra, together with other thorium experiments and supporting quantum chemical calculations, will be presented.

