

## SUB-DOPPLER SLIT JET DISCHARGE SPECTROSCOPY OF JET COOLED POLYACETYLENES: THE ANTI-SYMMETRIC CH STRETCH MODE OF TRIACETYLENE

CHIH-HSUAN CHANG, MELANIE A. ROBERTS, and DAVID J. NESBITT, *JILA, National Institute of Standards and Technology University of Colorado, and Department of Chemistry and Biochemistry University of Colorado at Boulder, Colorado 80309.*

Growth of polyacetylenic molecules in acetylene flames is thought to play a central role in combustion chemistry and formation of soot, as well as the chemistry of gas clouds in the interstellar medium. In this talk, we present results from first sub-Doppler, high resolution infrared spectroscopic studies on triacetylene.<sup>a,b</sup> In particular, we explore the fundamental anti-symmetric CH stretching mode ( $\nu_5$ ) of jet-cooled triacetylene in a pulsed slit discharge, where the *in-situ* synthesis arises from a discharge of trace(0.1-1%) acetylene/rare gas mixtures followed by rapid CCH + HCCH chemistry in the supersonic expansion environment. The band origin of this mode is determined to be  $3329.0544(2) \text{ cm}^{-1}$ . At high resolution, a series of avoided energy level crossings arising from rotational perturbations are observed and ascribed to perpendicular Coriolis mixing with a near degenerate manifold of  $\Pi$  vibrational symmetry. The energy level patterns are successfully analyzed to reveal spectroscopic constants and Coriolis coupling matrix elements for the perturbing manifold. In addition, a weak  $\Pi$ - $\Pi$  hot band progression due to thermal population in the slit jet is observed and assigned.

---

<sup>a</sup>D. McNaughton and D. N. Bruget, *J. Mol. Spectrosc.* 150, 620 (1991)

<sup>b</sup>K. Matsumura, K. Kawaguchi, D. McNaughton, and D. N. Bruget, *J. Mol. Spectrosc.* 158, 489 (1993)