

VIBRONIC SPECTROSCOPY OF $\text{MgCH}_3 \tilde{A}^2E \leftarrow \tilde{X}^2A_1$ TRANSITION

ALDO P. SALZBERG, BRIAN E. APPLGATE, TERRY A. MILLER, *Laser Spectroscopy Facility, Department of Chemistry, The Ohio State University, Columbus, OH 43210.*

We report the laser excitation and dispersed fluorescence spectra of the $\text{MgCH}_3 \tilde{A}^2E \leftarrow \tilde{X}^2A_1$ transition. Improvements to the experiment have allowed the observation of important vibronic features not previously observed. The radical was produced in a free jet expansion from the simultaneous photolysis of $\text{Hg}(\text{CH}_3)_2$ and the laser vaporization of solid Mg. The use of $\text{Hg}(\text{CH}_3)_2$ as a precursor substantially improves production of the radical. A 0.1 cm^{-1} -resolution laser was used to obtain the partially rotationally resolved LIF spectra of the 6_0^1 , and 2_0^1 transitions. Analysis of these spectra, along with the previously reported^a 0_0^0 band yield structural parameters which provide information about the radical's \tilde{A}^2E state. Besides the usual rotational constants, we have obtained information about the Jahn-Teller effect, which is dynamic and relatively weak. We have also obtained the values of the Coriolis coupling and the spin-orbit coupling constants. The fluorescence lifetime was found to be 60(7) ns for the vibrationless level of the \tilde{A} state and less than 15 ns for transitions involving vibrational excitation. The dispersed fluorescence spectra of the jet-cooled MgCH_3 was analyzed, and all observed 2A_1 state vibrational features were assigned.

^aR. Rubino, J. Williamson, and T. A. Miller, J. Chem. Phys. 103(14) (1995).