

HIGH RESOLUTION ELECTRONIC SPECTROSCOPY OF INDOLE AND INDOLE(H₂O) IN A MOLECULAR BEAM: EVIDENCE FOR ¹L_α- AND ¹L_β-STATE MIXING.^a

JOCHEN KÜPPER, *Heinrich-Heine-Universität, Institut für Physikalische Chemie und Elektrochemie I, 40225 Düsseldorf, Germany*; TIMOTHY M. KORTER, and DAVID W. PRATT, *University of Pittsburgh, Department of Chemistry, Pittsburgh, PA 15260, USA*.

The electronic spectrum of indole and its sensitivity to the surrounding environment continues to attract much attention.^b We report here studies of the rotationally resolved electronic spectra of several vibronic bands of indole and indole(H₂O) in a molecular beam, up to an energy of 990 cm⁻¹ above the electronic origin of the ¹L_β state at 35231.525 cm⁻¹ (≈284 nm). We find, as did Berden, et al.^c for the 0₀⁰ band, that the vibronic bands exhibit “anomalous” line intensities, evidencing axis tilting on S₁ excitation. The degree of tilt is different for different bands, showing that the S₁ ← S₀ optical transition moment orientations are also different. We derive from careful analyses of these data information about ¹L_α-¹L_β state mixing in the isolated molecule and its dependence on displacements along different vibrational coordinates. We also show that complexation of indole with one water molecule significantly influences the degree of this mixing.

^aWork supported by NSF(CHE-9617208)

^bY. Huang and M. Sulkes, *J. Phys. Chem.* 100(1996),16479 and references therein.

^cG. Berden, W.L. Meerts, and E. Jalviste, *J. Chem. Phys.* 103(1995), 9596