

## INTERNAL STATE DISTRIBUTIONS OF PRODUCT OH AND NO FROM PHOTODISSOCIATION OF NITRIC ACID AT 193 NM

G. H. LEU, I. C. CHEN, *Department of Chemistry, National Tsing Hua University, Hsinchu, Taiwan, ROC 30043.*

The distributions of vibrational, rotational, spin-orbit and lambda -doublet states of nascent NO and OH are obtained using the A-X transition of both OH and NO via the technique of laser-induced fluorescence. NO is produced from the secondary dissociation of fragment NO<sub>2</sub> from nitric acid after photolysis at 193 nm. About 3% of available energy is distributed to rotation of OH fragment; no vibrational excited OH was observed. It was found that about 87.7% of excess energy to be directed to the internal energy of fragment NO<sub>2</sub>. According to the measurement of internal state distributions of secondary product NO, the partition of energy in NO X<sup>2</sup>Π is calculated to be 840 cm<sup>-1</sup> in vibration and 1410 cm<sup>-1</sup> in rotation. The spin-orbit state <sup>2</sup>Π<sub>1/2</sub> of NO is populated about twice of that in <sup>2</sup>Π<sub>3/2</sub> for the vibrational levels v=0,1,2. The measured populations of vibrational states v=0,1,2 are 0.75, 0.19 and 0.07, respectively ; these populations agree with result calculated with the statistical model, prior theory. The experimental data indicate that although NO<sub>2</sub> photochemical from nitric acid , was proposed to be in an electronically excited state, the state distribution of NO from unstable NO<sub>2</sub> agree with those from NO<sub>2</sub> excited with monochromatic light in the UV-vis range after average for the broad distribution of internal energy. Hence this unknown electronic state of NO<sub>2</sub> is expected to be coupled to the ground electronic surface then leading to dissociation.