

LARGE-AMPLITUDE BENDING DYNAMICS OF ACETYLENE

MATTHEW P. JACOBSON, JONATHAN P. O'BRIEN, ROBERT J. SILBEY, and ROBERT W. FIELD, *Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139.*

We have investigated the large-amplitude bending dynamics of acetylene, in its ground electronic state, using an effective Hamiltonian model that reproduces all relevant experimental data (84 vibrational levels), up to $15,000 \text{ cm}^{-1}$ in internal energy, with 1.4 cm^{-1} accuracy (1σ). This investigation has been made possible by a numerical pattern recognition analysis of our dispersed fluorescence (DF) data set for the acetylene $\tilde{A}^1A_u \rightarrow \tilde{X}^1\Sigma_g^+$ system, which includes DF spectra recorded from five different vibrational levels of the \tilde{A}^1A_u state. Through this pattern recognition analysis, polyad quantum numbers have been assigned to observed transitions in the DF spectra up to $E_{vib} = 15,000 \text{ cm}^{-1}$. Our analysis of the “pure bending” polyads, which involve excitation exclusively in the *trans* and *cis* bending modes, has revealed a rich, but in many ways, surprisingly simple, dynamics at high internal energy ($> 10,000 \text{ cm}^{-1}$). Among the conclusions of this analysis is that, in many ways, the observed bending dynamics is somewhat simpler at $15,000 \text{ cm}^{-1}$ than it is at $10,000 \text{ cm}^{-1}$; this rather surprising result can be explained in terms of qualitative changes in the structures of the pure bending polyads as a function of increasing internal energy. In addition, the eigenfunctions of the effective Hamiltonian at high internal energy are classifiable in terms of “local bending” and “librational” motions; this observation implies that a localized model of the bending dynamics is more appropriate at high internal energy than a normal model model, and we have derived analytical expressions for converting between these two representations of the dynamics.