

THE PRODUCTION AND INFRARED SPECTRUM OF A PHOTODIMER OF TRIFLUORONITROSOMETHANE (CF₃NO) ISOLATED IN INERT GAS MATRICES

MEGAN MORGAN, *Titleist Foot Joy World Wide, Fairhaven, MA 02719*; and C. A. BAUMANN, *Department of Chemistry, University of Scranton, Scranton, PA 18510-4626*.

We report the results of laser photolysis ($\lambda = 670, 633$ nm) of CF₃NO trapped in solid argon, krypton, xenon, nitrogen and oxygen. Infrared spectroscopy was used to characterize the photoproducts and monitor the progress of the photolysis. In the gas phase photolysis, the parent molecule undergoes fragmentation upon S₁ excitation and the nascent fragments react with another monomer to produce the N-nitroamine, (CF₃)₂NONO, previously observed by Mason^a. Our condensed phase spectra indicate the formation of a different photoproduct, an azo or azoxy compound, with no apparent presence of the nitroamine, and traces of CF₃ and NO in only the most dilute matrices. The quantum efficiency of this reaction exhibits a dependence on the matrix:reagent ratio. This concentration dependence is consistent with a second photolytic pathway where the S₁ monomer reacts with a ground state monomer in an adjacent matrix site before fragmentation can occur.

^aJ. Mason, J. Chem. Soc. 1963, 4531.