

SPEED-DEPENDENT EFFECTS ON THE LINE SHAPES OF ARGON-BROADENED HF

A. S. PINE, *Alpine Technologies, 14401 Poplar Hill Road, Germantown, MD 20874.*

Asymmetric line shapes have been observed for infrared transitions of Ar-broadened HF^a and were fit to Dicke-narrowed profiles with an empirical partial correlation between velocity- and state-changing collisions. That analysis yielded measured broadening and shifting coefficients in excellent agreement with the thermally-averaged values calculated from exact quantum close-coupled scattering cross-section based on a realistic van der Waals potential^b. However, some anomalies in the sense of the asymmetries and the magnitude and J dependence of the velocity-changing rates remained, which subsequently were addressed classically by Joubert et al.^c and quantum mechanically by Demeio et al.^d. Here, we reexamine those line shapes with the correlated speed-dependent strong-collision Dicke-narrowed profiles described above using the reliable energy-dependent cross-sections of Green and Hutson. For HF/Ar below one atmosphere, Dicke narrowing dominates the speed dependence.

^aA. Pine, J. Chem. Phys. 101, 3444 (1994).

^bS. Green and J. Hutson, J. Chem. Phys. 100, 891 (1994).

^cP. Joubert, J. Bonamy, D. Robert, J. Domenech and D. Bermejo, JQSRT (to be published).

^dL. Demeio, S. Green and L. Monchick, J. Chem. Phys. 102, 9160 (1995).