

## IR-CRLAS OF SMALL BIOMOLECULES

C. J. CHAPO, J. B. PAUL, K. ROTH, and R. J. SAYKALLY, *Department of Chemistry, UC Berkeley, Berkeley, CA 94720.*

Infrared cavity ringdown laser absorption spectroscopy (IR-CRLAS) is a novel technique which has been used to study a wide variety of systems.<sup>a,b</sup> Currently, we report the application of this method to study hydration of small biomolecules. We have recorded mid-infrared absorption spectra for several jet-cooled amino acids, including valine and arginine. We have also collected spectra of (water)<sub>n</sub>-biomolecule clusters for these systems. In addition, we have recorded spectra in the Amide I region (1600–1700 cm<sup>-1</sup>) for jet-cooled GlyGly, which has allowed us to begin examination of peptide backbone conformation issues. Along with *ab initio* frequency calculations, these results provide insight in answering questions such as the gas-phase conformation distribution and the zwitterionic state of these systems.

---

<sup>a</sup>J. J. Scherer, D. Voelkel, D. J. Rakestraw, J. B. Paul, C. P. Collier, R. J. Saykally, and A. O'Keefe. *Chem. Phys. Lett.* **245**, 273 (1995).

<sup>b</sup>J. B. Paul, C. P. Collier, R. J. Saykally, J. J. Scherer, and A. O'Keefe. *J. Chem. Phys. A* **101**, 5211 (1997).