

HIGH PRECISION MEASUREMENT OF C-O STRETCHING BAND OF CH₃OH USING CO₂ LASER SIDEBANDS

Z. D. SUN, F. MATSUSHIMA, S. TSUNEKAWA, and K. TAKAGI, *Department of Physics, Toyama University, Toyama 930-8555, Japan.*

Microwave sidebands of CO₂ laser lines have been used as an infrared power source to measure the sub-Doppler vibration-rotation frequencies of CH₃OH in the C-O stretching band precisely. More than one hundred frequencies of CH₃OH have been measured. Over thirty groups of unresolved lines in the Fourier-transform spectrum have been resolved. Combined with our previous infrared-microwave double resonance results^a and the ground state ($\nu_{co} = 0$) term values given in literature^b, many term values of CH₃OH in the $\nu_{co} = 1$ stretching states have been refined. For A species in $\nu_{co} = 1$, 22 asymmetry splittings for various K in $\nu_t = 0$ and 1 have been observed for the first time. The present results provide precision data to refine the molecular constants in the first excited C-O stretching state of CH₃OH.

^aZ. D. Sun, S. Ogura, F. Matsushima, S. Tsunekawa and K. Takagi, paper RA08 at the "53"^d Ohio State University International Symposium on Molecular Spectroscopy", Columbus, Ohio (1998).

^bG. Moruzzi, B. P. Winnewisser, M. Winnewisser, I. Mukhopadhyay, and F. Strumia, *Microwave, infrared and laser transitions of methanol*, (CRC Press, Inc. 1995).