

THE LOW-LYING CCN BENDING MODE OF THE CLOSE-TO-LINEAR MOLECULE NCCNO

HOLGER LICHAU, MICHAEL LOCK, MANFRED WINNEWISSER, *Physikalisch-Chemisches Institut, Justus-Liebig-Universität, D-35392 Giessen, Germany*; STEPHEN C. ROSS, *Department of Physics, University of New Brunswick, Fredericton NB E3B 5A3, Canada*.

The band system of the low-lying CCN bending mode ν_7 of the pseudohalofulminate NCCNO around 85 cm^{-1} has been recorded by high resolution FTIR spectroscopy to supplement our previous investigation of the pure rotational spectrum, which had revealed some slight, but significant indications of quasilinearity^a. Besides rovibrational transitions of the fundamental band, rovibrational transitions belonging to the first, second, third, and fourth hot band could be assigned. Using a conventional linear-molecule type Hamiltonian^b, these rovibrational transitions have been analyzed together with the pure rotational transitions. The results of this analysis confirm a moderate anharmonicity in the CCN bending potential. To quantify this anharmonicity, the combined data have been subjected to a *semirigid bender* analysis, which resulted in a rather flat bending potential with a significant quartic contribution.

^aH. Lichau, B. P. Winnewisser, M. Winnewisser, C. W. Gillies, and J. Z. Gillies, *53rd Int. Symposium*, talk TH09 (1998)

^bK. M. T. Yamada, F. W. Birss, and M. R. Aliev, *J. Mol. Spectrosc.* **112**, 347–356 (1985)